Accepted Manuscript

Title: A case report on the successful perioperative management of hepatectomy for hepatocellular carcinoma in a patient with von Willebrand disease

Authors: Koki Sato, Shintaro Kuroda, Tsuyoshi Kobayashi, Seiichi Shimizu, Masahiro Ohira, Hiroyuki Tahara, Kentaro Ide, Teruhisa Fujii, Hideki Ohdan



S2210-2612(18)30067-1 https://doi.org/10.1016/j.ijscr.2018.02.033 USCR 3044

To appear in:

Received date:	1-12-2017
Revised date:	23-1-2018
Accepted date:	22-2-2018

Please cite this article as: Sato Koki, Kuroda Shintaro, Kobayashi Tsuyoshi, Shimizu Seiichi, Ohira Masahiro, Tahara Hiroyuki, Ide Kentaro, Fujii Teruhisa, Ohdan Hideki.A case report on the successful perioperative management of hepatectomy for hepatocellular carcinoma in a patient with von Willebrand disease.*International Journal of Surgery Case Reports* https://doi.org/10.1016/j.ijscr.2018.02.033

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

A case report on the successful perioperative management of hepatectomy for hepatocellular carcinoma in a patient with von Willebrand disease

Koki Sato¹, Shintaro Kuroda¹*, Tsuyoshi Kobayashi¹, Seiichi Shimizu¹, Masahiro Ohira¹, Hiroyuki Tahara¹, Kentaro Ide¹, Teruhisa Fujii², Hideki Ohdan¹

¹Department of Gastroenterological and Transplant Surgery, Hiroshima University Hospital, 1-2-3, Kasumi, Minami-ku, Hiroshima, 734-8551, Japan.

²Division of Blood Transfusion Services, Hiroshima University Hospital, 1-2-3, Kasumi, Minami-ku, Hiroshima, 734-8551, Japan.

*Corresponding author:

Shintaro Kuroda

Highlights

- Although von Willebrand disease (VWD) is a common inherited bleeding disease, only 1 case of hepatectomy in a patient with VWD has been described.
- Perioperative hemostatic management is important especially in patients with VWD undergoing surgery because of the dysfunction of the platelet and destabilization of the blood clotting factor VIII.
- The patient was successfully treated by administering factor VIII/von Willebrand factor concentrate and by measuring activated partial thromboplastin time as an index for perioperative hemostatic management.

ABSTRACT

INTRODUCTION: Although von Willebrand disease (VWD) is a common inherited

Download English Version:

https://daneshyari.com/en/article/8832804

Download Persian Version:

https://daneshyari.com/article/8832804

Daneshyari.com