




Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Challenging case of horseshoe kidney double fracture

Francesco Cortese^a, Pietro Fransvea^{d,*}, Roberto Marcello^b, Alessandra Saputelli^a,
Luca Lepre^a, Aldo Giofrè^a, Gabriele Sganga^c

^a Emergency Surgery and Trauma Care Unit, St. Filippo Neri's Hospital, Rome, Italy

^b Department of Radiology, Vascular and Interventional Radiology Unit, St. Filippo Neri's Hospital, Rome, Italy

^c Surgical Clinics, General and Transplantation Surgery Unit, Sacred Heart Catholic University, A. Gemelli Teaching Hospital, Rome, Italy

^d Faculty of Medicine and Psychology, "Sapienza" University of Rome, St. Andrea's Hospital, Italy

ARTICLE INFO

Article history:

Received 28 June 2016

Received in revised form 7 August 2017

Accepted 8 August 2017

Available online 10 October 2017

Keywords:

Renal trauma

Horseshoe kidney

Renal anatomy

ABSTRACT

INTRODUCTION: Renal injuries occur in 10% of blunt abdominal traumas, 7% of these occur in kidneys with congenital or acquired disorders. Trauma of horseshoe kidney is an uncommon finding.

PRESENTATION OF A CASE: We present the case of 31 year-old caucasian man with no remarkable personal records, who was brought to our Trauma Unit soon after being involved in a motorcycle collision. A Contrast Enhanced – Multi Detector Computed Tomography (ce-MDCT) revealed a double disconnection of a horseshoe kidney. The patient was not aware of bearing such abnormality.

DISCUSSION: Trauma of horseshoe kidney is an uncommon finding. The abdominal ce-MDCT scan is the diagnostic tool of choice since the renal anatomy, injury grading and vascular or urinary tract abnormalities are well depicted and easily identified. The conservative management of these injuries is associated with a lower rate of nephrectomies and kidney failure while selective *trans*-catheter renal embolization is a challenging treatment option. However surgery can be a treatment of choice and should be aimed to preserve renal function. Conclusion: the interest in our case lies in the rarity and particular anatomical aspect of such injuries and the implication related to its management in an emergency setting.

© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Renal trauma is a well-known accepted cause of significant morbidity and mortality at trauma centers across the globe, despite kidneys are located in a relatively protected retroperitoneal position. Renal injuries occur in 10% of blunt abdominal traumas, 7% of these occurring in kidneys with congenital or acquired disorders. Horseshoe kidney is a congenital disorder of the kidney characterized by the fusion of both kidney lower poles arising at the time of renal spin rotation around the main axis, typically appearing between week 4 and 6 of pregnancy. The two kidneys are often joined by a fibrous or parenchymatous isthmus connecting the two renal poles. These anomalous kidneys may have normal function although they show important anatomical changes in their vascular network and sometimes serious urinary tract abnormalities. Horseshoe kidneys occur in 1 out of 400 people, it is more common in males (ratio 2:1) and it is sometimes associated with other different congenital disorders. Approximately 60% of the horseshoe kidney patients are asymptomatic and diagnosis is obtained

as an incidental finding. In other cases, the most common symptom is related to recurring renal stones, UTI or hydronephrosis due to urinary tract obstruction. The increase of motor vehicle accidents and high speed collisions during the last 10 years, as well as the industrial and sport collisions and the soar of street violence and aggressiveness, increased renal trauma incidence in severe abdominal traumas. The particular anatomical aspect of the horseshoe kidney leads to an extreme vulnerability of such congenital condition in case of abdominal trauma; therefore, the correct and quick detection of this abnormality is needed in order to carry out a prompt and adequate treatment.

2. Case report

A white, 31 Y old male with no remarkable personal records, was brought to our Trauma Unit soon after he was involved in a motorcycle collision. His conditions were critical with hypovolemic shock due to haemorrhage. Blood pressure was 80 over 50 mmHg, heart rate was 140 bpm with a GCS being 13. Lab tests revealed hemoglobin value of 14,5 g/dl, hematocrit of 42,3%, WBC count 9.700/mm³, platelets count of 200.000/mm³, creatinine of 1,2 mg/dl, urea of 25 mg/dl and prothrombin of 82%. He also presented with left humeral and peroneal fracture. Primary survey was carried out according to ATLS[®] approach with good response and he was then rushed to undergo total body CT scan. A Contrast

* Corresponding author at: Faculty of Medicine and Psychology, University of Rome "La Sapienza", St. Andrea, Hospital, ViaDi Grottarossa, 1035-39, 00189 Rome, Italy.

E-mail address: pietro.fransvea@gmail.com (P. Fransvea).



Fig. 1. Ce-MDCT of abdomen showing large left perirenal hematoma and active extravasation of contrast media consistent with active bleeding.

Enhanced – Multi Detector Computed Tomography (ce- MDCT) revealed a double disconnection of a horseshoe kidney with a massive retroperitoneal haematoma. The patient was not aware of bearing such abnormality. Trauma injuries were considered of grade IV according to the American Association for the Surgery with an ISS 16 of Trauma classification (Fig. 1). No other abdominal injuries were detected. In order to attend a conservative treatment

the patient was sent to the Interventional Radiology Unit (IRU). The angiography performed confirmed a bilateral kidney transection with a retroperitoneal haematoma suspected evolving, with a selective peripheral/distal embolization resulting impossible due to either the anatomical abnormality, either the kidney malpositioning as compressed by the hematoma. A bilateral embolization of both renal arteries had to be carried out as a selective treatment was not possible. A *trans*-catheter embolization was not indicated because of the high risk chronic kidney failure. Also, a new CT scan revealed an increase in the hematoma volume indicating a high risk of imminent rupture (Fig. 2). Hemoglobin level dropped to 7, therefore the patient was taken immediately to the OR to undergo surgical laparotomy. The intraoperative findings included a quote of free abdominal fluid (400cc) and injury-free intraperitoneal organs. Nevertheless, the presence of a large fissuring retroperitoneal hematoma was confirmed. To achieve a successful retroperitoneal space exposure synchronous Cattel-Mattox (Left-to-right medial visceral rotation) and Cattel-Braasch (Right-to-left medial visceral rotation) maneuvers were performed. A complete bilateral disconnection of the middle segment of the horseshoe kidney was observed. Sutures of the injured renal segments were achieved without the need of a ureteral stenting deployment (Fig. 3). Three abdominal drainages were then placed, two perinephric and one in the pelvis. Two unit of packed red blood cell (PRC) were transfused. A temporary abdominal wall closure was carried out in order to prevent the onset of an abdominal compartment syndrome. Bone injuries were treated conservatively at the end of the abdominal surgery. Post – operative renal function was normal. On post-op day 14 the patient was discharged from hospital



Fig. 2. Abdominal aorta angiography and right and left renal artery selective angiography. Horseshoe kidney tear and discontinuation is very well depicted. Extravasation of contrast media on rupture sites is also shown.




Fig. 3. Intraoperative findings. Right and left horseshoes kidney upper poles sutured. Note the left ureter well depicted in the middle of the figure.

Download English Version:

<https://daneshyari.com/en/article/8833023>

Download Persian Version:

<https://daneshyari.com/article/8833023>

[Daneshyari.com](https://daneshyari.com)