

Reverse extra-anatomic aortic arch debranching procedure allowing thoracic endovascular aortic repair of a chronic ascending aortic aneurysm

Ludovic Canaud, MD, PhD, Bernard Albat, MD, PhD, Kheira Hireche, MD, Aurelien Hostalrich, MD, PhD, Pierre Alric, MD, PhD, and Thomas Gandet, MD, Montpellier, France

ABSTRACT

A 79-year-old woman was admitted with a large chronic dissecting ascending aortic aneurysm starting 5 mm distal to the ostia of the left coronary artery and ending immediately proximal to the innominate artery. A reverse extra-anatomic aortic arch debranching procedure was performed. During the same operative time, through a transapical approach, a thoracic stent graft was deployed with the proximal landing zone just distal to the coronary ostia and the distal landing zone excluding the origin of the left common carotid artery. The postoperative course was uneventful. Computed tomography at 12 months documented patent extra-anatomic aortic arch debranching and no evidence of endoleak. (*J Vasc Surg Cases and Innovative Techniques* 2018;4:102-5.)

Keywords: Ascending aorta; Thoracic aorta; Aortic dissection; Stent graft

Type A aortic dissection is a cardiovascular catastrophe associated with high mortality. Medical therapy is associated with poor outcome, with a mortality rate of around 60%.¹ Open surgical repair is the first-line therapy for type A aortic dissection. However, a mortality rate of up to 60%² has been reported in high-risk patients. A less invasive approach would be highly beneficial in high-risk patients.

Currently, stent graft placement in the ascending aorta for the treatment of pseudoaneurysms, intramural hematoma, and even type A dissections has been described in anecdotal reports.^{3,4} We report a “reverse” extra-anatomic debranching to provide a suitable landing zone for thoracic endovascular aortic repair (TEVAR). The patient has consented to the publication of this article.

CASE REPORT

The patient, a 79-year-old woman with a past medical history of nonischemic cardiomyopathy and severe chronic obstructive pulmonary disease, presented to the emergency department with chest pain and shortness of breath. Computed tomography (CT) showed a large chronic dissecting ascending aortic aneurysm 6.9 cm in anteroposterior diameter (Figs 1 and 2; Video).

From the Department of Thoracic and Cardiovascular Surgery, Hôpital A de Villeneuve.

Author conflict of interest: none.

Correspondence: Ludovic Canaud, MD, PhD, Service de Chirurgie Vasculaire et Thoracique, Hôpital A de Villeneuve, 191 av Doyen Gaston Giraud, 34090 Montpellier, France (e-mail: ludoviccanaud@hotmail.com).

The editors and reviewers of this article have no relevant financial relationships to disclose per the Journal policy that requires reviewers to decline review of any manuscript for which they may have a conflict of interest.

2468-4287

© 2018 The Author(s). Published by Elsevier Inc. on behalf of Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.jvscit.2018.02.004>

This patient was unlikely to survive open ascending arch repair because of her cardiac morbidity and severe chronic obstructive pulmonary disease. Emergent transthoracic echocardiography showed a dilated ascending aorta with an intimal flap, an ejection fraction of 40%, and diastolic dysfunction. Pulmonary evaluation performed 1 year ago found a postbronchodilator forced expiratory volume in 1 second of 75% of predicted value.

The aneurysm started 5 mm distal to the ostia of the left coronary artery and ended immediately proximal to the brachiocephalic trunk. The aorta above the aortic valve measured 31 mm, whereas the distal landing zone at the left subclavian was only 24 mm in diameter. A reverse arch debranching procedure was performed. This was composed of the following: left subclavian artery to right axillary bypass (retropharyngeal route); sequential reimplantation of the left and right common carotid arteries into the left subclavian artery to right axillary bypass; and right and left common carotid artery and right subclavian artery ligation. Ligation of the innominate artery was not accessible by a cervical approach. Bypass was performed using an 8-mm Dacron graft (Fig 3). During the same operative time, a transapical approach was performed. A left lateral thoracotomy in the fifth intercostal space (5-cm incision) was used to expose the apex of the heart. Two standard purse-string sutures were placed to close the hole at the end of the procedure, and an echocardiographically guided puncture was performed. A Valiant (Medtronic, Santa Rosa, Calif) stent graft (34 × 100 mm) was chosen. The graft was deployed under fluoroscopic control with the proximal landing zone just distal to the coronary ostia and the distal landing zone excluding the origin of the left common carotid artery. The proximal and distal end of the stent graft was precisely adjusted and deployed under rapid ventricular pacing. Completion angiography demonstrated patency of the coronary ostia and the left subclavian artery without endoleak. Balloon dilation was not performed.




Fig 1. Computed tomography (CT) showed a large dissecting ascending aortic aneurysm (6.9 cm). The aneurysm started 5 mm distal to the ostia of the left coronary artery and ended immediately proximal to the brachiocephalic trunk.




Fig 2. Contrast-enhanced axial computed tomography (CT) image demonstrates an intimal flap that separates the two channels in the ascending aorta.

After surgery, the patient was extubated within 24 hours. A CT scan confirmed satisfactory exclusion of the ascending aortic aneurysm with no endoleak. The patient made a successful recovery and was discharged home on the fifth postoperative day. CT at 12 months documented a patent stent graft and extra-anatomic aortic arch debranching and no evidence of endoleak, stent graft expansion, and aneurysm shrinkage (Fig 4).

DISCUSSION

This report demonstrates a case of chronic type A aortic dissection successfully treated by TEVAR. It illustrates the utility of reverse extra-anatomic aortic arch debranching and the transapical approach for precise deployment of stent grafts in the ascending aorta.

Emergency surgical intervention remains the “gold standard” treatment of type A aortic dissection and intramural hematoma, yet up to 30% of patients with this disease are unable to undergo an open repair.⁵ In those




Fig 3. Drawing of the reverse extra-anatomic aortic arch debranching and thoracic endovascular aortic repair (TEVAR) of a chronic ascending aortic aneurysm.

patients unfit for open repair, endovascular therapy has emerged as a possible alternative.

The anatomic and physiologic challenges to endovascular therapy of the ascending aorta remain formidable. They include proximal graft fixation close to the aortic valve and coronary ostia and a distal landing zone that may impinge on the innominate artery.

The proximal entry tear was located at the distal part of the ascending aorta at least 6 cm distal to the left coronary ostia but was associated with a retrograde extension of the dissection ending 5 mm distal to the left coronary ostia. We thought that even landing in a proximal dissected neck, the closure of the entry tear could seal the chronic dissecting aneurysm.

CT imaging studies have reported that TEVAR is anatomically feasible in about 30% of patients with type A dissection with a proximal entry tear in the ascending aorta.⁶

Extra-anatomic bypass has previously been described through left lateral thoracotomy from the descending thoracic aorta.⁷ This approach has the advantage of providing a long landing zone for the stent graft but at the cost of invasive thoracotomy and thoracic aorta clamping. This case report illustrates the feasibility of a reverse aortic arch debranching procedure to provide a distal suitable landing zone. The main advantages are to provide debranching from a cervical approach without aortic clamping.

Another challenge is that the hemodynamic forces in the ascending aorta resist accurate graft deployment. Rapid pacing helps improve stent positioning during

Download English Version:

<https://daneshyari.com/en/article/8835884>

Download Persian Version:

<https://daneshyari.com/article/8835884>

[Daneshyari.com](https://daneshyari.com)