Title: The iron pro-chelator BHAPI attenuates glutamate-induced oxidative stress via Wnt-β/Catenin Pathway in HT22 cells

Authors: Fei Gao, Bao Wang, Tao Chang, Min Li, Wei Fang, Zhi-hong Li, Li Gao

PII: S0361-9230(18)30212-0
DOI: https://doi.org/10.1016/j.brainresbull.2018.03.011
Reference: BRB 9397

To appear in: *Brain Research Bulletin*

Received date: 20-3-2018
Accepted date: 22-3-2018

Please cite this article as: Fei Gao, Bao Wang, Tao Chang, Min Li, Wei Fang, Zhi-hong Li, Li Gao, The iron pro-chelator BHAPI attenuates glutamate-induced oxidative stress via Wnt-β/Catenin Pathway in HT22 cells, Brain Research Bulletin https://doi.org/10.1016/j.brainresbull.2018.03.011

The iron pro-chelator BHAPI attenuates glutamate-induced oxidative stress via Wnt-β/Catenin Pathway in HT22 cells

Fei Gao[a,1], Bao Wang[a,1], Tao Chang[a], Min Li[a], Wei Fang[a], Zhi-hong Li[a], Li Gao[a*]

[a] Department of Neurosurgery, Tangdu Hospital, Fourth Military Medical University, Xi'an, Shaanxi 710038, China

[1] These authors contributed equally to this work.

[*]Correspondence to:

Dr. Li Gao,

Department of Neurosurgery, Tangdu Hospital,

Fourth Military Medical University,

Xi'an, Shaanxi Province 710038, China

Tel: 0086-29-84717556

Fax: 0086-29-84717556

Email: gaoli089@163.com

Abstract

Disturbances in intracellular iron homeostasis are associated with brain damage under various neuropathological conditions. However, exposure of neuronal cells to classical iron chelators could interfere with physiological iron functions in the brain.