Accepted Manuscript

Title: A Novel Method for Oxygen Glucose Deprivation Model in Organotypic Spinal Cord Slices

Authors: Jing-Jie Liu, Xiao-yan Ding, Li Xiang, Feng Zhao, Sheng-Li Huang



PII:S0361-9230(17)30061-8DOI:https://doi.org/10.1016/j.brainresbull.2017.10.010Reference:BRB 9312To appear in:Brain Research BulletinReceived date:5-2-2017Pavised date:10.10.2017

 Revised date:
 10-10-2017

 Accepted date:
 12-10-2017

Please cite this article as: Jing-Jie Liu, Xiao-yan Ding, Li Xiang, Feng Zhao, Sheng-Li Huang, A Novel Method for Oxygen Glucose Deprivation Model in Organotypic Spinal Cord Slices, Brain Research Bulletin https://doi.org/10.1016/j.brainresbull.2017.10.010

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

A Novel Method for Oxygen Glucose Deprivation Model in Organotypic Spinal Cord Slices

Jing-Jie Liu^a, Xiao-yan Ding^b, Li Xiang^a, Feng Zhao^c, Sheng-Li Huang^{c,*}

^aDepartment of Neurology, the Second Affiliated Hospital, College of Medicine, Xi'an Jiaotong University, Xi'an 710004, China

^bDepartment of Ophthalmology, Xi'an Third Hospital, Xi'an 710018, China.

^cDepartment of Orthopedics, the Second Affiliated Hospital, College of Medicine, Xi'an Jiaotong University, Xi'an 710004, China.

*Corresponding author: Sheng-Li **Huang**, Department of Orthopedics, the Second Affiliated Hospital, College of Medicine, Xi'an Jiaotong University, No. 157 Xiwulu, Xi'an 710004, China

Tel: + 86 298 767 9621; e-mail address: huangshengli@xjtu.edu.cn.

Highlights

- This study established a model to closely mimic spinal cord hypoxic-ischemic injury.
- OGD model is obtained from long-term cultured organotypic spinal cord slices.
- For OGD model, 60 minutes was the optimal time.

Download English Version:

https://daneshyari.com/en/article/8839070

Download Persian Version:

https://daneshyari.com/article/8839070

Daneshyari.com