



Review Article

Preventing synaptic deficits in Alzheimer's disease by inhibiting tumor necrosis factor alpha signaling

Chelsea Cavanagh^a, Tak Pan Wong^{a,b,*}^a Douglas Mental Health University Institute, Montreal, QC, Canada^b Department of Psychiatry, McGill University, Montreal, QC, Canada

ARTICLE INFO

Article history:

Received 20 November 2017

Accepted 31 January 2018

Keywords:

Amyloid precursor protein

Hippocampus

Neuroinflammation

Prevention

TgCRND8 mice

Tumor necrosis factor alpha

XPro1595

ABSTRACT

The characterization of preclinical stages of Alzheimer's disease (AD) would provide a therapeutic window for prevention. One of the challenges of developing preventive therapy for AD is to identify early biomarkers for intervention studies. We have recently shown that in the TgCRND8 transgenic AD mouse model, increased hippocampal levels of the pro-inflammatory cytokine tumor necrosis factor alpha (TNF α) and enhanced excitatory synaptic transmission were early-onset changes that occurred weeks before amyloid plaque formation. Inhibiting TNF α before plaque formation not only normalized excitatory synaptic function, but also prevented the impairment of synaptic function 4 months later. In this review paper, we will examine the potential contributions of TNF α to the alteration of brain function in preclinical AD. The prospective use of TNF α inhibitors for preventing AD will be discussed.

© 2018 The Authors. Published by Elsevier Ltd on behalf of International Brain Research Organization. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

Acknowledgements.....	20
References.....	20

Alzheimer's disease (AD) is a neurodegenerative disorder and the most common cause of dementia, which affects close to 50 million people worldwide (AD-International, 2015). While the prevalence of AD is expected to be tripled by the mid-21st century, effective AD treatment is lacking. Findings from at-risk genetic cohorts or older asymptomatic people have revealed a long pathophysiological process that starts years, if not decades, before the onset of clinical AD. Several preclinical AD stages before the onset of overt cognitive decline have been proposed and are currently characterized by studying changes in imaging modalities and biomarkers such as amyloid and tau levels in cerebrospinal fluid (Sperling et al., 2011). While the accumulated brain damage in patients with AD dementia is likely to be irreversible, preclinical AD could be a time window for preventive treatment. Indeed, while increased AD risk has been associated with mid-life factors, such as hypertension, physical inactivity and poor education (Barnes and

Yaffe, 2011), clinical trials have revealed some protective effects of antihypertensive therapy, exercise, and cognitive training (Andrieu et al., 2015). An effective preventive treatment could have a significant impact on the economic burden of AD to society, since delaying the onset age of AD by 5 years has been estimated to reduce the incidence of AD by 50% (Brookmeyer et al., 1998). A challenge for developing a preventive treatment for AD is to characterize and validate earlier biomarkers for intervention studies.

Neuroinflammation could be an early stage biomarker for preclinical AD. While the idea that anti-inflammatory treatment underlies the low prevalence of AD in rheumatic arthritis patients has been around for over 25 years (McGeer et al., 1990), the role of neuroinflammation remains unclear in the transition from asymptomatic preclinical stages to AD dementia. Pro-inflammatory agents like cytokines have been implicated in the pathophysiology of AD. For example, interferon- γ (IFN- γ) mRNA expression is elevated in the Tg2576 mouse model of AD at early stages of amyloid pathology (Abbas et al., 2002). In addition, TNF α along with monocyte chemoattractant protein were also shown to be increased before amyloid plaque deposition in the entorhinal cortex of 3xTgAD mice, another mouse model of AD (Janelins et al.,

* Corresponding author at: Douglas Mental Health University Institute, 6875 LaSalle Blvd Montreal, QC, H4H 1R3, Canada.
E-mail address: takpan.wong@mcgill.ca (T.P. Wong).

2005). Evidence of neuroinflammation is also present in patients with mild cognitive impairment who have increased caspase-1 (Heneka et al., 2013). The TgCRND8 mouse model that expresses the human amyloid precursor protein (APP) gene with both the Indiana (V717F) and Swedish (K670N/M671L) mutations, which leads to an overproduction of amyloid- β (A β). Using this model, we found that the levels of the pro-inflammatory cytokine, tumor necrosis factor α (TNF α) rise in 1-month old mice, 2 months before the onset of amyloid pathology and cognitive impairment (Cavanagh et al., 2013), which becomes prevalent at 3 months of age in this model (Chishti et al., 2001). This early-stage increase in TNF α levels can have a significant impact on brain functions. TNF α enhances neuronal excitability by inserting the AMPA subtype of glutamate receptors and removing GABA_A receptors from synapses (Stellwagen and Malenka, 2006). Notably, enhanced hippocampal activation has been observed in cognitively normal carriers of apolipoprotein E4 (ApoE4) (Nichols et al., 2012) or presenilin 1 (PS1) gene mutations (Reiman et al., 2012), and these individuals also have an increased risk of developing dementia. Enhanced hippocampal activation may reflect an increase in activity at the synaptic level. Therefore, we hypothesize that increased TNF α signaling at an early stage may lead to synaptic hyperexcitability, which could precipitate synaptic and cognitive pathology in AD.

In a recent study (Cavanagh et al., 2016), we examined the contribution of enhanced levels of TNF α on functional properties of glutamate synapses in the hippocampus of pre-plaque (1-month old) TgCRND8 mice. We observed stronger glutamatergic transmission in the hippocampus of pre-plaque TgCRND8 mice than in control, nontransgenic littermates. At this young age, TgCRND8 mice also exhibited higher hippocampal synaptic (long-term potentiation) and cognitive function (passive avoidance) than control mice (Cavanagh et al., 2016). Moreover, enhanced hippocampal synaptic and cognitive functions were normalized by the TNF α inhibitor, XPro1595 (Cavanagh et al., 2016). This novel biologic is a dominant-negative version of TNF α that can form inactive trimers with endogenous soluble TNF α to thereby prevent TNF α receptor 1 (TNFR1) activation (Steed et al., 2003), which mediates signaling that is often pathogenic and can lead to cell death (Van Hauwermeiren et al., 2011). These findings support a role of TNF α in enhancing hippocampal synaptic function before amyloid plaque formation. To determine the impact of the early-stage increase in TNF α on the formation of synaptic impairments that have been observed in post-plaque 6–12 month old TgCRND8 mice (Kimura et al., 2012), we administered XPro1595 in 1-month old TgCRND8 mice through a subcutaneously implanted osmotic pump for 4 weeks. After pump removal, mice were left undisturbed until they became 6-month old. Compared with control mice, 6-month old TgCRND8 mice exhibited a decrease in glutamatergic transmission in the hippocampus. Decreased synaptic function however was rescued in XPro1595-treated TgCRND8 mice (Cavanagh et al., 2016). To our knowledge, this was the first study to show that inhibiting TNF α signaling before amyloid plaque formation prevented the development of synaptic pathology at a later stage in an AD mouse model (Cavanagh et al., 2016). Inhibiting TNF α signaling in 1-month old TgCRND8 mice may normalize the increase in synaptic transmission that is also found at this stage. This increase in synaptic transmission in TgCRND8 mice could be detrimental to synaptic function at later stages, since normalizing the increase by inhibiting TNF α prevented the development of synaptic deficits at advanced stages.

Findings from Cavanagh and colleagues support a hypothesis that increased hippocampal activity is an early-stage change in AD. Previous findings in other APP mouse models have revealed increased neuronal activation near diffuse plaques (Busche et al., 2008) and aberrant electroencephalogram activity (Palop et al., 2007). These functional changes could be related to the increased

risk of seizures in AD patients (Pandis and Scarmeas, 2012), especially in cohorts of early-onset and familial AD (Cabrejo et al., 2006; Scarmeas et al., 2009). In human studies, increased hippocampal activity has been observed in asymptomatic carriers of AD-related genes (Nichols et al., 2012; Reiman et al., 2012). Furthermore, in a longitudinal study, the levels of cortical activity in ApoE4 carriers correlated with future degree of cognitive impairment (Bookheimer et al., 2000), suggesting a causal relationship between a preclinical increase in brain activity and cognitive decline. The increase in synaptic function we observed in young TgCRND8 mice could have a significant impact on the hippocampal network, since measuring hippocampal oscillations of pre-plaque TgCRND8 mice has revealed a decrease in theta-gamma cross frequency coupling (Goutagny et al., 2013). While mechanisms underlying the preclinical increase in brain activity in AD remain unclear, they could be related to alterations in inhibitory synapses. The reduction of GABAergic synaptic function has been suggested to underlie enhanced neuronal activation near amyloid plaques in the frontal cortex (Busche et al., 2008). Moreover, decreased GABAergic transmission resulting from decreased expression of the voltage-gated sodium channel subunit Nav1.1 has been detected in the hAPP-J20 mouse model as well as in AD patients (Verret et al., 2012) and GABAergic neurons have been shown to be vulnerable to A β toxicity (Krantic et al., 2012). Although we observed increased evoked excitatory synaptic function in the hippocampal CA1 region of pre-plaque TgCRND8 mice, we found no change in miniature excitatory synaptic transmission (Cavanagh et al., 2016). These findings suggest that alterations at the network level, may be related to a reduction in inhibitory synaptic inputs that could underlie the enhanced evoked synaptic responses we observed in these mice. However, changes in inhibitory function could be compensated after enhanced excitability. In the J20 AD mouse model, seizure activity in the hippocampal region was associated with the sprouting of inhibitory inputs in the dentate gyrus (Palop et al., 2007). Further studies using young asymptomatic APP mouse models are needed to reveal synaptic mechanisms that are responsible for causing neuronal hyperexcitability in APP mouse models.

Before amyloid plaque formation, an increase in inflammatory mediators, such as activated microglia (Wright et al., 2013) and cytokines (Goutagny et al., 2013; Hanzel et al., 2014; Janelins et al., 2005), has been observed in cortical and hippocampal regions of APP rodent models. Data from our intervention study suggest that TNF α inhibitors may be useful in preventing synaptic deficits in AD. TNF α can enhance neuronal excitability by increasing and decreasing the expression of sodium (Chen et al., 2015) and potassium channels (Diem et al., 2001), respectively. Excitatory synaptic transmission can also be enhanced by TNF α through insertion of calcium permeable GluA2-lacking AMPA receptors (Leonoudakis et al., 2008), which could be mediated by TNFR1 on astrocytes (Habbas et al., 2015). TNF α also enhances the insertion (Wheeler et al., 2009) and calcium influx (Jara et al., 2007) of the NMDA subtype of glutamate receptors. Finally, TNF α can facilitate glutamate release from astrocytes (Santello et al., 2011). Alternatively, activation of neuronal TNFR1 can lead to the endocytosis of GABA_A receptors (Pribiag and Stellwagen, 2013). While the pro-excitation effects of TNF α are important for synaptic scaling (Stellwagen and Malenka, 2006), an activity-dependent physiological process for refining neuronal networks, the chronic increase in TNF α we observed in TgCRND8 mice (Cavanagh et al., 2013) could result in hyperexcitability and enhanced sensitivity to excitotoxicity. Increased levels of pro-inflammatory markers like other cytokines (e.g. IFN- γ (Abbas et al., 2002), interleukin1 β , interleukin 6 (Heneka et al., 2005)) and the complement system (Hong et al., 2016) have been reported in pre-plaque APP mouse models. Therefore, future studies should find out whether an early-stage increase in these

Download English Version:

<https://daneshyari.com/en/article/8839101>

Download Persian Version:

<https://daneshyari.com/article/8839101>

[Daneshyari.com](https://daneshyari.com)