
Accepted Manuscript

Title: Comparison among ultrasonic, electrical apparatus, and toxic chemicals for vestibular lesion in mice

Authors: Yusuke Yamaoka, Chikara Abe, Hironobu Morita



Please cite this article as: Yamaoka Yusuke. Abe Chikara. Morita Hironobu.Comparison among ultrasonic, electrical apparatus, and toxic chemicals for vestibular lesion in mice. Journal of Neuroscience Methods https://doi.org/10.1016/j.jneumeth.2017.11.021

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Comparison among ultrasonic, electrical apparatus, and toxic chemicals for vestibular lesion in mice

By

Yusuke Yamaoka, Chikara Abe, and Hironobu Morita

Department of Physiology, Gifu University Graduate School of Medicine, Gifu, Japan

Article type: Research paper

Word count of manuscript: 6036

Total number of figures: 8

Author responsible for correspondence:

Chikara Abe, DDS & PhD

Department of Physiology, Gifu University Graduate School of Medicine

1-1 Yanagido, Gifu, 501-1194, Japan

Tel: +81-58-230-6300

Fax: +81-58-230-6302

Email: chikara@gifu-u.ac.jp

Highlights

- Vestibular lesion induced using an ultrasonicator shows complete deletion of vestibular function without non-specific damage.
- Vestibular lesion can be induced chemically in C57BL/6J but not in ICR mice; however, the degree of disruption of hair cells was lower than that using an ultrasonicator or electrical apparatus.

Download English Version:

https://daneshyari.com/en/article/8840453

Download Persian Version:

https://daneshyari.com/article/8840453

Daneshyari.com