

Accepted Manuscript

Entrainment of the circadian clock in neural stem cells by epidermal growth factor is closely associated with ERK1/2-mediated induction of multiple clock-related genes

Asuka Mogi, Ryo Yomoda, Syunya Kimura, Chisato Tsushima, Jun Takouda, Miho Sawauchi, Tomoko Maekawa, Hidenobu Ohta, Satoshi Nishino, Masatake Kurita, Nariyasu Mano, Noriko Osumi, Takahiro Moriya

PII: S0306-4522(18)30170-2

DOI: <https://doi.org/10.1016/j.neuroscience.2018.02.045>

Reference: NSC 18335


To appear in: *Neuroscience*

Received Date: 10 October 2017

Accepted Date: 27 February 2018

Please cite this article as: A. Mogi, R. Yomoda, S. Kimura, C. Tsushima, J. Takouda, M. Sawauchi, T. Maekawa, H. Ohta, S. Nishino, M. Kurita, N. Mano, N. Osumi, T. Moriya, Entrainment of the circadian clock in neural stem cells by epidermal growth factor is closely associated with ERK1/2-mediated induction of multiple clock-related genes, *Neuroscience* (2018), doi: <https://doi.org/10.1016/j.neuroscience.2018.02.045>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Entrainment of the circadian clock in neural stem cells by epidermal growth factor is closely associated with ERK1/2-mediated induction of multiple clock-related genes

Asuka Mogi^{a†}, Ryo Yomoda^{ab†}, Syunya Kimura^{ab}, Chisato Tsushima^{ab}, Jun Takouda^a, Miho Sawauchi^{ab}, Tomoko Maekawa^a, Hidenobu Ohta^{cd}, Satoshi Nishino^{abe}, Masatake Kurita^{abe}, Nariyasu Mano^f, Noriko Osumi^g, and Takahiro Moriya^{abf}

^aDepartment of Cellular Signaling, ^bLaboratory of Pharmacotherapy of Life-Style Related Diseases, Graduate School of Pharmaceutical Sciences, Tohoku University, Aramaki, Sendai, Japan.

^cDepartment of Psychophysiology, National Institute of Mental Health, National Center of Neurology and Psychiatry, 4-1-1 Ogawa-higashi-cho, Kodaira, Tokyo 187-8553, Japan

^dDepartment of Psychiatry, Asai Hospital, 38-1 Togane, Chiba 283-0062, Japan

^eNatori Station Front Clinic, 428-1 Hachiman Tegurada, Natori, Miyagi 981-1231, Japan

^fFaculty of Pharmaceutical Sciences, Tohoku University; Department of Pharmaceutical Sciences, Tohoku University Hospital, 1-1 Seiryomachi, Aoba-ku, Sendai, Japan

^gDepartment of Developmental Neuroscience, Center for Neuroscience, United Centers for Advanced Research and Translational Medicine, Tohoku University School of Medicine, Sendai, Japan

* Corresponding author: Takahiro Moriya, Ph.D., Department of Cellular Signaling, Graduate School of Pharmaceutical Sciences, Tohoku University, Aramaki, Aoba-ku, Sendai 980-8578, Japan. (Phone: +81-22-795-3843; Fax: +81-22-795-3847; e-mail: moriya@m.tohoku.ac.jp)

[†] Equally contributed to this study.

Download English Version:

<https://daneshyari.com/en/article/8840759>

Download Persian Version:

<https://daneshyari.com/article/8840759>

[Daneshyari.com](https://daneshyari.com)