Accepted Manuscript



Title: Diagnostic model for attention-deficit hyperactivity disorder based on interregional morphological connectivity

Authors: Xun-Heng Wang, Yun Jiao, Lihua Li

PII: S0304-3940(18)30502-0
DOI: https://doi.org/10.1016/j.neulet.2018.07.029
Reference: NSL 33719

To appear in: *Neuroscience Letters*

Received date: 1-3-2018
Revised date: 22-6-2018
Accepted date: 19-7-2018

Please cite this article as: Wang X-Heng, Jiao Y, Li L, Diagnostic model for attention-deficit hyperactivity disorder based on interregional morphological connectivity, *Neuroscience Letters* (2018), https://doi.org/10.1016/j.neulet.2018.07.029

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Diagnostic model for attention-deficit hyperactivity disorder based on interregional morphological connectivity

Xun-Heng Wang[a*], Yun Jiao[b], and Lihua Li[a*]

[a] College of Life Information Science and Instrument Engineering, Hangzhou Dianzi University, Hangzhou, 310018, China

[b] Jiangsu Key Laboratory of Molecular and Functional Imaging, Department of Radiology, Zhongda Hospital, Medical School of Southeast University, Nanjing, 210009, China

Corresponding

Xun-Heng Wang, Ph.D. (Email: xhwang@hdu.edu.cn, Phone: 0571-87713533, Fax: 0571-87713528)

Lihua Li, Ph.D. (Email: lilh@hdu.edu.cn, Phone: 0571-86878587, Fax: 0571-87713528)

Highlights

- Individual morphological connectivity contains high-order cortical information.
- Interregional features outperform regional measures in diagnostic model.
- Discriminative morphological connectivity can be discovered by linear SVM.
- Individual morphological connectivity correlates to clinical symptoms.
- Individual morphological connectivity can be a biomarker for ADHD.

ACCEPTED MANUSCRIPT

Download English Version:

https://daneshyari.com/en/article/8841312

Download Persian Version:

https://daneshyari.com/article/8841312

Daneshyari.com