

Title: Wider range of allodynia in a rat model of repeated dural nociception compared with infraorbital nerve chronic constriction injury

Authors: Guanqun Hu, Mingjie Zhang, Min Su, Qing Zhang, Hangfei Wu, Xiaolin Wang, Zhao Dong, Shengyuan Yu

PII: S0304-3940(17)31021-2
DOI: https://doi.org/10.1016/j.neulet.2017.12.048
Reference: NSL 33320

To appear in: *Neuroscience Letters*

Received date: 2-11-2017
Revised date: 16-12-2017
Accepted date: 22-12-2017

Please cite this article as: Guanqun Hu, Mingjie Zhang, Min Su, Qing Zhang, Hangfei Wu, Xiaolin Wang, Zhao Dong, Shengyuan Yu, Wider range of allodynia in a rat model of repeated dural nociception compared with infraorbital nerve chronic constriction injury, Neuroscience Letters https://doi.org/10.1016/j.neulet.2017.12.048

Title: Wider range of allodynia in a rat model of repeated dural nociception compared with infraorbital nerve chronic constriction injury

Authors: Guanqun Hu[1,2], Mingjie Zhang[1,*], Min Su[1,2], Qing Zhang[1,2], Hangfei Wu[1,2], Xiaolin Wang[1], Zhao Dong[1], Shengyuan Yu[1,*]

1. Department of Neurology, Chinese PLA General Hospital, Beijing 100853, China

2. School of Medicine, Nankai University, Tianjin, 300071, China

*Corresponding Author(s):

The corresponding authors are Dr. Shengyuan Yu and Dr. Mingjie Zhang.

Shengyuan Yu: Department of Neurology, Chinese PLA General Hospital, Beijing 100853, China

E-mail: yusy1963@126.com

Mingjie Zhang: Department of Neurology, Chinese PLA General Hospital, Beijing 100853, China

E-mail: mjzhangnk@163.com

* These two authors contributed equally to this work.

Article Highlights

- Cephalic and extra-cephalic allodynia exist in the interictal period in IS rats.
- Wider range of allodynia in the IS model compared with the IoN-CCI model.
- IS and IoN-CCI can be seen as suitable animal models of CM and TN, respectively.