

Accepted Manuscript



A 3' UTR-derived non-coding RNA RibS increases expression of *cfa* and promotes biofilm formation of *Salmonella enterica* serovar Typhi

Xin Zhao, Rui Liu, Hao Tang, George Osei-Adjei, Shungao Xu, Ying Zhang, Xinxiang Huang

PII: S0923-2508(18)30063-9

DOI: [10.1016/j.resmic.2018.04.007](https://doi.org/10.1016/j.resmic.2018.04.007)

Reference: RESMIC 3653

To appear in: *Research in Microbiology*

Received Date: 29 December 2017

Revised Date: 12 April 2018

Accepted Date: 12 April 2018

Please cite this article as: X. Zhao, R. Liu, H. Tang, G. Osei-Adjei, S. Xu, Y. Zhang, X. Huang, A 3' UTR-derived non-coding RNA RibS increases expression of *cfa* and promotes biofilm formation of *Salmonella enterica* serovar Typhi, *Research in Microbiology* (2018), doi: 10.1016/j.resmic.2018.04.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

For publication

A 3' UTR-derived non-coding RNA RibS increases expression of *cfa* and promotes biofilm formation of *Salmonella enterica* serovar Typhi

Xin Zhao, Rui Liu, Hao Tang, George Osei-Adjei, Shungao Xu, Ying Zhang, Xinxiang

Huang*

Department of Biochemistry and Molecular biology, School of medicine, Jiangsu University,

Zhenjiang, Jiangsu, 212013, China^a

* Correspondence and reprints huxinx@ujs.edu.cn

Xin Zhao, 453391197@qq.com; Rui Liu, 970483911@qq.com;

Hao Tang, thly191012@126.com; George Osei-Adjei, geoadjei@hotmail.com;

Shungao Xu, 1000007337@ujs.edu.cn; Ying Zhang, zhangyingzj85@163.com;

Xinxiang Huang, huxinx@ujs.edu.cn

Abstract

Bacterial non-coding RNAs (ncRNAs) are widely studied and found to play important roles in regulating various cellular processes. Recently, many ncRNAs have been discovered to be transcribed or processed from 3' untranslated regions (3' UTRs). Here we reported a novel 3' UTR-derived ncRNA, RibS, which could influence biofilm formation of *Salmonella*

Download English Version:

<https://daneshyari.com/en/article/8842840>

Download Persian Version:

<https://daneshyari.com/article/8842840>

[Daneshyari.com](https://daneshyari.com)