



## Original article

Ovarian development and histological observations of threatened dwarf snakehead fish, *Channa gachua* (Hamilton, 1822)James Milton<sup>a</sup>, Ajaz A. Bhat<sup>a</sup>, M.A. Haniffa<sup>a</sup>, Shaikh Althaf Hussain<sup>b,\*</sup>, Irfan A. Rather<sup>c,1</sup><sup>a</sup> Centre for Aquaculture Research and Extension (CARE), St. Xavier's College (Autonomous), Palayamkottai 627002, Tamil Nadu, India<sup>b</sup> Central Laboratory, Zoology Department, College of Science, King Saud University, PO Box 2455, Riyadh 11451, Saudi Arabia<sup>c</sup> Department of Applied Microbiology and Biotechnology, Yeungnam University, Gyeongsan, Gyeongbuk 712-749, South Korea

## ARTICLE INFO

## Article history:

Received 12 December 2016

Revised 16 January 2017

Accepted 13 March 2017

Available online xxxx

## Keywords:

Histology

Ovary

Gonadosomatic index

*Channa gachua*

## ABSTRACT

*Channa gachua* were monthly sampled throughout a year and the histological analysis of their ovaries was done to determine the changes occurring in ovarian development. Based on histological examination of the ovaries, the oogenic process of *C. gachua* undergoes distinct cyclic and seasonal morphological changes. Five different developmental stages were identified under three major categories: pre-spawning (immature, maturing, mature), spawning (ripe-running) and post-spawning (spent). The peak spawning period of *C. gachua* was noticed during December - February. The gonadosomatic index (GSI) and ova diameter ranged from 0.79 to 3.61% and 543–1123  $\mu\text{m}$  respectively. The highest mean GSI ( $3.61 \pm 0.16$ ) and oocyte diameter ( $1123 \pm 55 \mu\text{m}$ ) were observed in December indicating that during this month the gonadal development reached maturity.

© 2017 Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Murrels commonly called snakeheads due to the presence of large scales on their head belong to the family Channidae. *Channa gachua*, commonly called dwarf snakehead is an important food fish and is one of the most suitable channa species for aquarium due to its beautiful colouration and small size (Talwar and Jhingran, 1992). Even though it is widely distributed in Asia, it has declined drastically (vulnerable) in India (CAMP, 1998) and endangered in some Asian countries like Singapore (Lim and Ng, 1990). Although, this species is economically important, information on the reproductive physiology and histomorphological changes of ovary of *C. gachua* in captive conditions remains limited.

For proper management of aquaculture practices, a detailed study of gonad maturation is important, since such studies are aimed in understanding the annual changes of the population

(Thorpe et al., 1990; Jobling et al., 2002; Tomkiewicz et al., 2003; Shein et al., 2004). The reproductive cycles in teleost occur during a particular phase; some breed once in a year as annual breeders and others as monsoon breeders (Zuckerman, 1962).

Freshwater murrels being seasonal breeders exhibit clear changes in the gonads during breeding season (Marimuthu et al., 2001). Cyclic gonadal changes have been studied in a few *Channa* species viz. *Channa marulius* (Parameswaran and Murugesan, 1976), *C. punctata* (Srivastav and Srivastav, 1998) and *C. striata* (Al Mahmud et al., 2016).

Histological studies on fish reproduction to determine peak period of spawning and to understand effective methods for increasing efficiency of broodstock and ultimately increasing the fish production are prerequisite. The detailed information on changes in the ovaries of the dwarf snakehead during the reproductive cycle is important to provide useful data for the management of this species. Hence, this study was performed to examine the ovarian developmental stages of dwarf snakehead, *C. gachua* during its annual maturation cycle under captive conditions.

## 2. Materials and methods

## 2.1. Sample collection

The experimental fishes (12–20 cm in total length, 15–50 g in weight) were collected from Thamirabarani River (8.44°N,

Peer review under responsibility of King Saud University.



\* Corresponding author.

E-mail addresses: [althafbio@gmail.com](mailto:althafbio@gmail.com) (S.A. Hussain), [rather@ynu.ac.kr](mailto:rather@ynu.ac.kr) (I.A. Rather).<sup>1</sup> Co-corresponding author.<http://dx.doi.org/10.1016/j.sjbs.2017.03.011>

1319-562X/© 2017 Production and hosting by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).Please cite this article in press as: Milton, J., et al. Ovarian development and histological observations of threatened dwarf snakehead fish, *Channa gachua* (Hamilton, 1822). Saudi Journal of Biological Sciences (2017), <http://dx.doi.org/10.1016/j.sjbs.2017.03.011>

77.44°E) by a cast net during the months of December 2008–February 2009. The fishes were acclimatized to captive conditions by maintaining them in earthen ponds (3 × 3 × 1 m) filled with chlorine free tap water (dissolved oxygen 5.2 mg l<sup>-1</sup>, temperature 28 ± 2 °C, and pH 6.4–7.1) at Center for Aquaculture Research and Extension (CARE) Aquafarm. The pond was provided with plenty of aquatic plants (*Hydrilla verticillata*). The fishes were fed on formulated diets (60% chicken intestine, 17% ground nut oil cake, 11% rice bran, 10% tapioca, and 2% vitamin and mineral mixture) following Haniffa et al. (1999), twice daily (morning and evening) until satiation.

## 2.2. Gonadosomatic index

Ovary samples were obtained from 5 female dwarf snakehead specimens every month captured using drag net from April 2009 to March 2010. Fish were measured (nearest 0.1 cm) and weighed (nearest 0.1 g) and were dissected to remove the ovaries. Ovarian samples were weighed (nearest 0.1 g) and fixed in 10% buffered formalin prior to histological analysis. Gonadosomatic index (GSI) was calculated using the following formula:

$$GSI = \frac{GW}{TW} \times 100$$

where

GW = Gonad weight

TW = Total body weight.

## 2.3. Histology

Gonad sections were collected from the mid-part of the ovary monthly for histological analysis. These sections were fixed in 10% buffered formalin for further laboratory analysis. After washing in running water, the samples were dehydrated in an ascending series of ethanol (70%, 90% and absolute ethanol) and clarified with xylene. The samples embedded in paraffin blocks were trimmed and 5 μm thick sections taken were stained using haematoxylin and eosin followed by periodic acid Schiff (PAS) reaction. The maturity stages of ovary of *C. gachua* were determined using the modified gonad maturity scale developed by Arockiaraj et al. (2004). The histological sections were examined under the light microscope (Nikon microscope - U III E-400 Eclipse). Diameter of eggs was measured using Magnus Pro Microscope Software with accuracy level of 0.01 mm.

One-way analysis of variance (ANOVA) was used to analyze the data, followed by comparison of means using Duncan's multiple range tests. Statistically significant differences were accepted at  $p < 0.05$ .

**Table 1**

Ovarian condition of different maturity stages and comparison with histological examination in *Channa gachua*.

Stage and period	Macroscopic appearance	Histological examination
I. Pre-spawning		
(a) Immature (June–July)	Ovary small, thin ribbon like transparent, whitish gray, granular in appearance occupying half of the abdominal body cavity, eggs very minute distinct under microscope	Monolayer follicle phase in primary oocyte development with stage I present, ovigerous lamellae from the tunica albuginea were evident (Fig. 3)
(b) Maturing (August–September)	Ovary straight; ova visible through the capsule; ova orange yellow in colour	Oogonia, chromatin nucleolar and perinuclear oocytes growing rapidly; cortical alveoli start to appear in few oocytes (Fig. 4)
(c) Mature (October–November)	Ovary increases in size; forms lobes and is the largest organ in the abdominal cavity; ova yellowish-orange in colour	Ovaries dominated by late perinuclear oocytes and primary yolk vesicle; few secondary yolk vesicle oocytes present (Figs. 5–7)
II. Spawning		
(d) Ripe-running (December–February)	Ovary fully distended and fills the abdominal cavity; oocyte orange yellow and easily shed on application of slight pressure on the ovary	Ovary dominated by tertiary oocytes; fully mature eggs with yolk globules; few previtellogenic stages begin to grow for the subsequent season (Figs. 8 and 9)
III. Post-spawning		
(e) Spent (March–May)	Ovary flaccid and often haemorrhagic if spawning was successful; few oocytes visible giving the ovary a speckled appearance	Post-ovulatory follicles, oocytes undergoing atresia and type I and II atretic oocytes. (Fig. 9)

## 3. Results and discussion

Stages of oocyte development similar to those described in most teleost fishes (Encina and Granado-Lorencio, 1997) were identified and described in *C. gachua*. The three maturity stages of oocyte development of *C. gachua* determined based on morphological features are described in Table 1. First stage was pre-spawning stage which was divided into three sub-stages viz, (a) immature, (b) maturing, (c) mature. Second stage was spawning or ripe-running stage and the third was post-spawning or spent stage. Mature ovaries (stage Ic) were frequently observed after October and were abundant in December to February signalling the period of spawning. Females with empty ovaries, spent fishes (stage IIIe) were seen between March and May indicating the recovery period. These different stages corresponded with those described macroscopically for *M. montanus* (Arockiaraj et al., 2004). However, microscopic examination of ovary sections reported by Al Mahmud et al. (2016) revealed four stages of maturity in striated snakehead, *Channa striata*. The female fish showed significant ( $P < 0.05$ ) weight changes in the ovaries corresponding to the three gametogenic stages (pre-spawning, spawning and post spawning) during different months of the year (Table 2).

To determine the GSI values, 60 female dwarf snakehead samples (weight range = 30–47 g) were used. The GSI of *C. gachua* was significantly different during the sampling months ( $P < 0.05$ ). The GSI was minimum in April ( $0.79 \pm 0.11\%$ ), began to increase in June and reached the stage of complete sexual maturity in December ( $3.61 \pm 0.16\%$ ) where the ovaries were ripe and mature (Table 2, Fig. 1). Our result showed highest GSI in winter months (December, January and February) which could be the possible spawning season of *C. gachua*. Moreover, the mean ova diameter gradually increases in size from March onwards and rapidly reaching the maximum size ( $1123 \pm 55 \mu\text{m}$ ) in December (Fig. 2). During the sampling months, the oocyte diameter was significantly different ( $P < 0.05$ ). The highest proportion of “spent” gonads was noticed in March - May, where the GSI values declined rapidly after spawning.

In contrast to our reports, Al Mahmud et al. (2016) reported the minimum GSI value during the month of September and higher values from April to July for female *Channa striata*. Similarly, highest values of GSI were recorded during April- July in *Channa bleheri* (Rinku et al., 2013) and during May - August in *C. punctatus* (Sunita et al., 2011; Lalta et al., 2011). The highest/lowest values of GSI were due to the active somatic energy accumulation/depletion (Encina and Granado-Lorencio, 1997). The variations are probably due to the fact that the changes in energy are usually more than the seasonal weight variations as was reported by Scott et al. (1980). In the current study, highest oocyte diameter was observed in December, indicating that the maturation of oocyte reached peak in this month.

Download English Version:

<https://daneshyari.com/en/article/8849998>

Download Persian Version:

<https://daneshyari.com/article/8849998>

[Daneshyari.com](https://daneshyari.com)