Accepted Manuscript

Environmental Health Risk Posed by Contamination of the Individual Water Wells

Justyna Kubicz, Adam Pawełczyk, Paweł Lochyński

PII: S0045-6535(18)31039-7

DOI: 10.1016/j.chemosphere.2018.05.182

Reference: CHEM 21514

To appear in: Chemosphere

Received Date: 29 March 2018

Accepted Date: 29 May 2018

Please cite this article as: Justyna Kubicz, Adam Pawełczyk, Paweł Lochyński, Environmental Health Risk Posed by Contamination of the Individual Water Wells, *Chemosphere* (2018), doi: 10.1016/j.chemosphere.2018.05.182

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

ENVIRONMENTAL HEALTH RISK POSED BY CONTAMINATION

1

2	OF THE INDIVIDUAL WATER WELLS
3	Justyna Kubicz ^a , Adam Pawełczyk ^b , Paweł Lochyński ^c
4	^a Wrocław University of Environmental and Life Sciences, Institute of Environmental Engineering
5	Norwida 25, 50-375 Wrocław, Poland
6	kubicz.justyna@upwr.edu.pl
7	
8	^b Wroclaw University of Science and Technology, Wyspianskiego27, 50-370 Wroclaw, Poland
9	(corresponding author)
10	adam.pawelczyk@pwr.edu.pl
11	phone: +48.71.3202930, +48.602587244, fax: +48.71.327280
12	
13	^c Wrocław University of Environmental and Life Sciences, Institute of Environmental Engineering
14	Norwida 25, 50-375 Wrocław, Poland
15	pawel.lochynski@upwr.edu.pl
16	
17	Abstract This paper presents the results of analyzes of water samples taken from individual
18	water wells intended for supplying households in non-urbanized areas with low population and
19	infrastructure density. Relatively high concentrations of harmful pollutants were found in
20	ground waters of the examined spots located in the suburban areas which may increase the risk
21	of health incidents in exposed populations. Based on long-term measurements carried out in
22	individual spots and toxicological data, the individual health risks related to carcinogenic
23	effects in humans were assessed, resulting from direct ingestion of such water and from dermal
24	contact. Non-carcinogenic hazard levels were determined, as well. In this work a novel
25	approach to health risk assessment was proposed. It consists of determining the risks that range

Download English Version:

https://daneshyari.com/en/article/8850733

Download Persian Version:

https://daneshyari.com/article/8850733

<u>Daneshyari.com</u>