## **Accepted Manuscript**

Phytoremediation Potential Of Poplar And Willow Species In Small Scale Constructed Wetland For Boron Removal

Kubilay Yıldırım, Gözde Çıtır Kasım

PII: S0045-6535(17)32004-0

DOI: 10.1016/j.chemosphere.2017.12.036

Reference: CHEM 20416

To appear in: Chemosphere

Received Date: 07 September 2017

Revised Date: 04 November 2017

Accepted Date: 06 December 2017

Please cite this article as: Kubilay Yıldırım, Gözde Çıtır Kasım, Phytoremediation Potential Of Poplar And Willow Species In Small Scale Constructed Wetland For Boron Removal, *Chemosphere* (2017), doi: 10.1016/j.chemosphere.2017.12.036

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



### **ACCEPTED MANUSCRIPT**

#### Research highlights

- Eight poplar and willow species tested in constructed wetland (CW) to investigate their Boron (B) phytoremediation ability from wastewater
- Poplar and willow species in CW removed 65-44% of total B in the wastewater
- Photostabilization (31-15%) was the major B removal pathway for some species in CW
- Some species could photoextract more 4000mg/kg B within their tissues
- B loaded wood and leaf materials could be used to produce wooden panels resistant to burn

#### Download English Version:

# https://daneshyari.com/en/article/8852505

Download Persian Version:

https://daneshyari.com/article/8852505

<u>Daneshyari.com</u>