

Locating emergent trees in a tropical rainforest using data from an Unmanned Aerial Vehicle (UAV)

Cici Alexander^{a,g,*}, Amanda H. Korstjens^a, Emma Hankinson^a, Graham Usher^b, Nathan Harrison^a, Matthew G. Nowak^{b,c}, Abdullah Abdullah^d, Serge A. Wich^{e,f}, Ross A. Hill^a

^a Bournemouth University, Department of Life and Environmental Sciences, Talbot Campus, Poole, Dorset, BH12 5BB, United Kingdom

^b The PanEco Foundation - Sumatran Orangutan Conservation Programme, Chileweg 5, Berg am Irchel 8415, Switzerland

^c Southern Illinois University, Department of Anthropology, 1000 Faner Drive, Carbondale, IL, 62901, USA

^d Syiah Kuala University, Department of Biology, Banda Aceh 23111, Indonesia

^e Liverpool John Moores University, School of Natural Sciences and Psychology, L33AF, Liverpool, United Kingdom

^f University of Amsterdam, Institute for Biodiversity and Ecosystem Dynamics, Sciencepark 904, Amsterdam 1098, Netherlands

^g Aarhus University, Aarhus Institute of Advanced Studies (AIAS), Høegh-Guldbergs Gade 6B, DK-8000 Aarhus C, Denmark

ARTICLE INFO

Keywords:

Habitat mapping
Drones
Point cloud
Sleeping trees
Conservation
Rainforest
Sumatra

ABSTRACT

Emergent trees, which are taller than surrounding trees with exposed crowns, provide crucial services to several rainforest species especially to endangered primates such as gibbons and siamangs (Hylobatidae). Hylobatids show a preference for emergent trees as sleeping sites and for vocal displays, however, they are under threat from both habitat modifications and the impacts of climate change. Traditional plot-based ground surveys have limitations in detecting and mapping emergent trees across a landscape, especially in dense tropical forests. In this study, a method is developed to detect emergent trees in a tropical rainforest in Sumatra, Indonesia, using a photogrammetric point cloud derived from RGB images collected using an Unmanned Aerial Vehicle (UAV). If a treetop, identified as a local maximum in a Digital Surface Model generated from the point cloud, was higher than the surrounding treetops (Trees_{EM}), and its crown was exposed above its neighbours (Trees_{SL}; assessed using slope and circularity measures), it was identified as an emergent tree, which might therefore be selected preferentially as a sleeping tree by hylobatids. A total of 54 out of 63 trees were classified as emergent by the developed algorithm and in the field. The algorithm is based on relative height rather than canopy height (due to a lack of terrain data in photogrammetric point clouds in a rainforest environment), which makes it equally applicable to photogrammetric and airborne laser scanning point cloud data.

1. Introduction

Non-human primates are an essential component of tropical biodiversity and they play important roles in forest regeneration and ecosystem health (Chapman et al., 2013). Arboreal primates spend a significant part of their days moving through the canopy, and about half of their life at sleeping sites, with most species rarely climbing down to the ground in suitable habitats with tall well-connected trees. Unlike larger apes such as orang-utans (*Pongo* spp.), smaller apes such as hylobatids do not build nests. Instead, hylobatids prefer to sleep in liana-free emergent trees with exposed crowns that have limited accessibility from surrounding trees, to avoid predators and provide a high vantage point (Anderson, 1998). Abundance of secure and stable sleeping sites, along with other factors, may be crucial for the survival of hylobatids, under the threats of increased deforestation and climate change

(Cheyne et al., 2012; Reichard, 1998).

Remote sensing has improved our understanding of the habitat preferences of birds and mammals (Goetz et al., 2007; Palminteri et al., 2012) by providing a continuous representation of the forest canopy. A limitation of ground based surveys is that data are collected only for small sample areas or plots. Furthermore, ground-based surveys in dense tropical forests are time-consuming, with complex multi-layered canopies and sometimes difficult terrain limiting visibility and access. Airborne Laser Scanner (ALS) data have been used to relate the presence and movement patterns of primates to forest structure, based on canopy height, closure and connectivity (Davies et al., 2017; McLean et al., 2016). ALS has distinct advantages over other remote sensing techniques in describing the three-dimensional structure of forests throughout their vertical profile, and capturing underlying terrain information. However, these data are still expensive to acquire, especially

* Corresponding author at: Aarhus Institute of Advanced Studies (AIAS), Aarhus University, Høegh-Guldbergs Gade 6B, DK-8000, Aarhus C, Denmark.
E-mail address: cici@aias.au.dk (C. Alexander).




Fig. 1. Estimated locations of potential sleeping trees (Trees_SL) overlaid on an ortho-photo mosaic of the study area; the area within the red square is shown in Fig. 2. Inset: Location of the study site (in red) in Sumatra (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

for small areas, such as mapping the territories of groups of primates.

Unmanned Aerial Vehicles (UAVs) are a low-cost alternative to manned aircraft for collecting data from small areas (Puliti et al., 2017), and UAV data have been used for rapid and efficient location of nests of chimpanzees (*Pan* spp.) and orang-utans (*Pongo* spp.) (van Andel et al., 2015; Wich et al., 2015). Photogrammetric point clouds on a forest canopy surface can be generated from an RGB camera mounted on a UAV. One of the main differences between photogrammetric and ALS point clouds, is the absence of points below dense forest canopy in the former. Unlike ALS, photogrammetric UAV point clouds are generated through image matching only on surfaces captured by the camera. This makes it very difficult to generate a reliable terrain model in dense forests from UAV data, which is essential for deriving canopy height (Dandois and Ellis, 2013; Lisein et al., 2013; Puliti et al., 2015; Tuominen et al., 2015).

Emergent trees are identified in the field based on their relative height from neighbouring trees, which could be estimated using UAV data, even in the absence of a terrain model. Although emergent trees provide essential services to a range of species such as langurs (*Presbytinae*), fruit bats (*Megachiroptera*) and eagles (*Nisaetus* spp.) in addition to hylobatids, and have been shown to be a major contributor to rainfall recycling (Holzman, 2009; Kunert et al., 2017), their detection, mapping and monitoring have been largely overlooked in earlier studies. The main aim of this study was therefore to assess the suitability of UAV point cloud data for locating emergent trees (and therefore potential sleeping trees for hylobatids) in a tropical rainforest in Northern Sumatra, Indonesia.

2. Study Area and Datasets

The study site is in Sikundur in the Leuser Ecosystem in Northern Sumatra, the only known place where three ape species, orang-utans (*Pongo abelii*), white-handed gibbons (*Hylobates lar*) and siamangs (*Symphalangus syndactylus*), still co-exist (Palombit, 1996). Airborne data from three flights were collected using a UAV system comprising a Skywalker UAV (1.7 m wingspan), fitted with an APM 2.6 autopilot module, RfD900 long-range telemetry and a GoPro Hero3 Black Edition

camera, between 22nd and 25th January 2015. The average flying altitude was 198 m above the launch location, covering an area of approximately 11.2 sq km, and generated 5400 images. An area of 6.5 sq km (centre: 98.07 °E; 3.96 °N) along the Besitang River, with known presence of gibbons and siamangs, was used as the study area.

3. Methods

3.1. Initial selection of treetops

An ortho-photo mosaic with a pixel size of 25 cm, a Digital Surface Model (DSM) with a grid size of 50 cm and a point cloud with an average density of 16.59 points m^{-2} , were generated from the UAV data using Structure from Motion (SfM) and photogrammetric algorithms implemented in Agisoft PhotoScan v1.3.0. The DSM was clipped to the study area and a slope raster was generated in ArcMap™ 10.1. Locations of tree tops were initially identified as grid cells in the DSM which were local maxima within a circular neighbourhood of 5-m radius (Trees_LM); a circular neighbourhood of 5 m identified most of the prominent canopy trees based on visual analysis.

3.2. Locations of emergent trees

Trees were selected as emergent trees if their treetops were the local maxima within a circular neighbourhood of 25-m radius and were at least 5 m taller than the surrounding treetops (Trees_EM). Since this forest has been selectively logged in the past, and very few trees in a similar study site in the region were found to have a crown radius larger than 12.5 m (Alexander et al., 2017), a neighbourhood radius of 25 m was considered to be adequate. Trees_EM was thus a subset of Trees_LM.

Sleeping trees of hylobatids have been observed to often have exposed crowns, with the trunk visible above the canopies of surrounding trees. The slope of the DSM represents the height difference between adjacent grid cells; a slope of 85° would correspond to an elevation difference of 5.72 m for a cell size of 50 cm. High slopes would also indicate less connectivity to the surrounding trees. The slope raster

Download English Version:

<https://daneshyari.com/en/article/8867757>

Download Persian Version:

<https://daneshyari.com/article/8867757>

[Daneshyari.com](https://daneshyari.com)