

Accepted Manuscript

Upper Pleistocene trace fossils from Ponta das Bicudas, Santiago, Cape Verde Islands: Systematics, taphonomy and palaeoenvironmental evolution

E. Mayoral, A. Santos, J.A. Gámez Vintaned, J. Ledesma-Vazquez, B.G. Baarli, M. Cachão, C.M. da Silva, M.E. Johnson


PII: S0031-0182(17)30760-5
DOI: doi:[10.1016/j.palaeo.2018.03.003](https://doi.org/10.1016/j.palaeo.2018.03.003)
Reference: PALAEO 8691

To appear in: *Palaeogeography, Palaeoclimatology, Palaeoecology*

Received date: 19 July 2017
Revised date: 2 March 2018
Accepted date: 2 March 2018

Please cite this article as: Mayoral, E., Santos, A., Gámez Vintaned, J.A., Ledesma-Vazquez, J., Baarli, B.G., Cachão, M., da Silva, C.M., Johnson, M.E., Upper Pleistocene trace fossils from Ponta das Bicudas, Santiago, Cape Verde Islands: Systematics, taphonomy and palaeoenvironmental evolution, *Palaeogeography, Palaeoclimatology, Palaeoecology* (2018), doi:[10.1016/j.palaeo.2018.03.003](https://doi.org/10.1016/j.palaeo.2018.03.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Upper Pleistocene trace fossils from Ponta das Bicudas, Santiago, Cape Verde Islands:
systematics, taphonomy and palaeoenvironmental evolution

E. Mayoral^{a*}, A. Santos^a, J.A. Gámez Vintaned^b, J. Ledesma-Vázquez^c, B.G. Baarli^d, M.
Cachão^e, C.M. da Silva^e, M.E. Johnson^d

^aDepartamento de Ciencias de la Tierra, Facultad de Ciencias Experimentales, Campus de El Carmen, Universidad de Huelva, Avda. 3 de marzo, s/n, 21071 Huelva, Spain, mayoral@uhu.es; asantos@dgyu.uhu.es

^bDepartment of Geosciences, Faculty of Sciences & Information Technology, Universiti Teknologi PETRONAS (UTP), 32610 Bandar Seri Iskandar (Tronoh), Perak Darul Ridzuan, Malaysia, jose.gamez@utp.edu.my

^cFacultad de Ciencias Marinas, Universidad Autónoma de Baja California, Ensenada, BC 22800, Mexico, ledesma@uabc.edu.mx

^dDepartment of Geosciences, Williams College, Williamstown, MA 01267, USA, gbaarli@williams.edu, markes.e.johnson@williams.edu

^eDepartamento de Geologia and Instituto Dom Luiz de Geociências, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal, mcachao@fc.ul.pt, paleo.carlos@fc.ul.pt

*Corresponding author. Phone: +003495921985; Fax: +0034 959 219440; E-mail address: mayoral@uhu.es (E. Mayoral)

ABSTRACT

Bioturbation structures from Ponta das Bicudas in Santiago Island, Cape Verde Archipelago are located in coarse-grained biocalcudite sandwiched between basaltic lavas of the Upper Pleistocene Assomada Formation. Four ichnospecies have been

Download English Version:

<https://daneshyari.com/en/article/8868243>

Download Persian Version:

<https://daneshyari.com/article/8868243>

[Daneshyari.com](https://daneshyari.com)