

Accepted Manuscript

Whole root system water conductance responds to both axial and radial traits and network topology over natural range of trait variation

Martin Bouda, Craig Brodersen, James Saiers

PII: S0022-5193(18)30357-6
DOI: [10.1016/j.jtbi.2018.07.033](https://doi.org/10.1016/j.jtbi.2018.07.033)
Reference: YJTBI 9558



To appear in: *Journal of Theoretical Biology*

Received date: 17 August 2017
Revised date: 30 May 2018
Accepted date: 24 July 2018

Please cite this article as: Martin Bouda, Craig Brodersen, James Saiers, Whole root system water conductance responds to both axial and radial traits and network topology over natural range of trait variation, *Journal of Theoretical Biology* (2018), doi: [10.1016/j.jtbi.2018.07.033](https://doi.org/10.1016/j.jtbi.2018.07.033)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- We extend the Landsberg and Fowkes (1978) model of water uptake to root networks.
- Dichotomous branching pattern is more efficient in water uptake than herringbone.
- Network topology affects uptake over reported ranges of hydraulic parameters.
- Total water uptake responded more to radial flow in lupin but axial flow in wheat.
- Axial flow limitation may affect monocot root habit evolution.

Download English Version:

<https://daneshyari.com/en/article/8876433>

Download Persian Version:

<https://daneshyari.com/article/8876433>

[Daneshyari.com](https://daneshyari.com)