Accepted Manuscript

Theory of turbid microalgae cultures.

Carlos Martínez, Francis Mairet, Olivier Bernard

PII:S0022-5193(18)30340-0DOI:10.1016/j.jtbi.2018.07.016Reference:YJTBI 9541

To appear in:

Journal of Theoretical Biology

Received date:13 February 2018Revised date:13 June 2018Accepted date:13 July 2018

Please cite this article as: Carlos Martínez, Francis Mairet, Olivier Bernard, Theory of turbid microalgae cultures., *Journal of Theoretical Biology* (2018), doi: 10.1016/j.jtbi.2018.07.016

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Highlights

- Average growth rate is determined by the maximum and minimum light intensities in the PBR.
- There exists an optical depth maximizing the average growth rate.
- There is an optimal incident light intensity depending on the turbidity of the culture.
- Incident light can inhibit the growth rate even in dense cultures.
- Conditions for the occurrence of Allee effect are determined.

1

Download English Version:

https://daneshyari.com/en/article/8876450

Download Persian Version:

https://daneshyari.com/article/8876450

Daneshyari.com