




REVISTA BRASILEIRA DE  
**Entomologia**  
A Journal on Insect Diversity and Evolution

[www.rbentomologia.com](http://www.rbentomologia.com)



Systematics, Morphology and Biogeography

## Redescription and first record in South America of *Neogriphoneura timida* Curran (Diptera, Lauxaniidae)

Ramon Luciano Mello<sup>a,\*</sup>, Thiago Mahlmann<sup>b</sup>, Vera Cristina Silva<sup>c</sup>

<sup>a</sup> Universidade Federal de Mato Grosso do Sul, Laboratório de Sistemática de Díptera, Campo Grande, MS, Brazil

<sup>b</sup> Instituto Nacional de Pesquisas da Amazônia, Manaus, AM, Brazil

<sup>c</sup> UNESP – Universidade Estadual Paulista, Faculdade de Ciências Agrárias e Veterinárias, Departamento de Morfologia e Fisiologia Animal, Jaboticabal, SP, Brazil

### ARTICLE INFO

#### Article history:

Received 4 March 2016

Accepted 17 October 2016

Available online xxx

Associate Editor: Marcia Souto Couri

#### Keywords:

Distribution

Schizophora

Sperm competition

Spermathecae

Spermatozoa

### ABSTRACT

The genus *Neogriphoneura* Malloch, 1924 is currently composed of 11 species with New World distribution. *Neogriphoneura timida* Curran, 1942 is recorded for the first time in South America, with occurrences in Brazil, Colombia and Trinidad and Tobago; and new Central American records are presented: Belize, Honduras and Mexico. Here we redescribe the species and present, for the first time, illustrations of the male terminalia and female spermathecae, and discuss the main diagnostic characteristics of external morphology. A brief discussion about the postcopulatory sexual selection in this species is proposed based on morphology of the spermathecae.

© 2016 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Introduction

Lauxaniidae is a large family of Schizophora flies comprised of over 2000 species in about 200 genera worldwide. In the Neotropical Region is composed by nearly 400 species in about 70 genera (Gaimari and Silva, 2010).

The genus *Neogriphoneura* was erected by Malloch (in Malloch and McAtee, 1924) to include the species *Sapromyza sordida* Wiedemann, 1830, originally recorded from West Indies. The genus *Rhabdolauxania* was described by Hendel (1925) for his two new species *R. laevifrons* from Peru, and *R. schnusei* (the type species) from Bolivia and Peru. *Neogriphoneura striatifrons* was later described by Hendel (1932) based on material from Bolivia. In a subsequent publication, Hendel (1933) described *Rhabdolauxania immaculata* from Paraguay. *Neogriphoneura timida* was described by Curran (1942) based on four males and six females from Panama; in addition describing two other new species: *N. striga* and *N. tertia* both based on material from Panama and Brazil.

After 66 years without any additional species or taxonomic modification, Mello and Silva (2008a) presented a taxonomic review of *Neogriphoneura*. In this review, they described the species *N. bispoi* from Brazil, *N. corrugata* from the British Virgin Islands, and

*N. pacata* from Bolivia; the genus *Rhabdolauxania* was synonymized under *Neogriphoneura*, and its three species were combined into *Neogriphoneura*; lectotypes for *R. immaculata* and *R. laevifrons* were designated; an identification key to the species was presented; and new occurrence records were registered for the following species: *N. sordida*, *N. striatifrons*, *N. striga*, *N. tertia*, and *N. timida*. According with this information, *Neogriphoneura* is comprised of the following 11 species: *N. bispoi*, *N. corrugata*, *N. immaculata*, *N. laevifrons*, *N. pacata*, *N. schnusei*, *N. sordida*, *N. striatifrons*, *N. striga*, *N. tertia*, and *N. timida*.

In this study, *N. timida* is redescribed, with photographs of external morphology, and illustrations of male terminalia, and female spermathecae; new records are added to its distribution, including the first record for the species in South America, in addition to new records within Central America and Mexico. The differentiated morphology of the spermathecae of this species, shared with two other congeners, *N. bispoi* and *N. corrugata*, is briefly discussed in the light of a possible correlated evolution with some male traits.

### Material and methods


The material analyzed in this study belongs to the following collections: American Museum of Natural History (AMNH), New York, USA; Coleção Zoológica da Universidade Federal de Mato Grosso do Sul (ZUFMS), Campo Grande, Brazil; Coleção Zoológica do Maranhão, Universidade Estadual do Maranhão (CZMA), Caxias,

\* Corresponding author.

E-mail: [ramon.mello@ufms.br](mailto:ramon.mello@ufms.br) (R.L. Mello).

<http://dx.doi.org/10.1016/j.rbe.2016.10.001>

0085-5626/© 2016 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



**Figs. 1–4.** *Neogriphoneura timida* Curran. (1) Habitus, lateral view (female); (2) habitus, dorsal view (female); (3) head, dorsal view (male); (4) wing (female). Scale: 2 mm.

Brazil; Instituto de Investigación de Recursos Biológicos Alexander von Humboldt (IAvH), Bogotá, Colombia; Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil; National Museum of Natural History (USNM), Washington, DC, USA; The Natural History Museum (BMNH), London, England.

The morphological terminology followed [Cumming and Wood \(2009\)](#). The dissection of the male terminalia follows the procedures presented in [Mello and Silva \(2008b\)](#).

### Taxonomy

*Neogriphoneura timida* Curran (Figs. 1–9)

*Neogriphoneura timida* Curran, 1942: 75. Type locality: Panama, Patilla Point. Holotype male, AMNH.

**Diagnosis:** This species can be separated from its congeners by the following combination of characters: frons unmarked; mesonotum with a central dark stripe anteriorly, and an oval spot on posterior margin extending onto scutellum; wing unspotted.

### Redescription

**Body** (Figs. 1 and 2): Yellow to yellowish brown or orange; length: 4–4.5 mm; wing: 3.2 mm.

**Head** (Figs. 1–3): Vertex rounded; ocellar tubercle brown, close to vertex, well developed; frons yellow, without dark markings, rectangular, wider than long, concave on middle part of anterior half; eye suboval, higher than long, posteroventral margin concave; face yellowish to light brown, unmarked, lower half slightly convex in profile; parafacial pale, whitish yellow; gena yellow; occiput and median occipital sclerite yellowish, unmarked; antenna yellow to yellowish brown, scape shorter than pedicel, first flagellomere two times longer than wide, arista about 2.5 times longer than the length of the antennal articles, black and plumose; palpus and labellum yellow to yellowish brown. Chaetotaxy: outer vertical

seta lateroconclinate, 1/3 length of inner vertical seta; inner vertical seta reclinate, postocellar seta convergent; ocellar seta diminutive; anterior orbital seta inclinate, slightly longer than reclinate posterior seta.

**Thorax** (Fig. 2): Mesonotum yellowish brown to orange, with a black stripe anteromedially and a rounded spot on posterior part, extending onto scutellum; scutellum flat, otherwise yellow; pleura yellowish. Chaetotaxy: 1 postpronotal seta; 0 + 3 dorsocentral setae; 1 prescutellar acrostichal seta; 1 presutural supra-alar seta; 1 postsutural supra-alar seta; 2 notopleural setae; postalar seta absent; 1 intra-alar seta; 1 proepisternal seta; 1 anepisternal seta; 1 katepisternal seta; 2 scutellar setae divergent. Legs: whitish yellow, tarsi yellow. Chaetotaxy: fore coxa with a transversal row of anterodorsal setae; fore femur with a longitudinal row of posterodorsal setae, a longitudinal row of 3 apical setae on ventral surface, and 2 apical setae on posterior surface; fore tibia with 1 preapical dorsal seta; mid coxa with a transversal row of setae on dorsal surface; mid femur with a longitudinal row of anterodorsal setae and 1 apical seta on posterior surface; mid tibia with 1 preapical dorsal seta and 1 strong tibial spur; hind femur with 2 longitudinal setae on apical margin of dorsal surface; hind tibia with 1 apical seta on dorsal surface. Wing (Fig. 4): hyaline; veins yellow; costa sapromyziform; R bare; r-m slightly basal to midpoint of discal medial cell; crossvein dm-cu at the midpoint of cell  $r_{4+5}$ . Halter yellowish brown to orange.

**Abdomen** (Fig. 2): Yellowish brown, some specimens with one oval dark spot medially on each of tergites 3–5. Male terminalia: Epandrial complex (Figs. 5 and 6): epandrium saddle-shape in posterior view (Fig. 5), lower and upper margins oval and straight respectively, in lateral view (Fig. 6), covered by large and medial setae; surstylus articulated to epandrium, conical in posterior view (Fig. 5), trapezoidal on lateral view (Fig. 6), lower margin covered by thin hairs in lateral view. Phallic complex (Figs. 7 and 8): phallopodeme tubular, well developed (Fig. 7); phallus cylindrical, folded in the middle, with sclerotized spines internally on apical region (Figs. 7 and 8); hypandrial arms developed, almost three

Download English Version:

<https://daneshyari.com/en/article/8877278>

Download Persian Version:

<https://daneshyari.com/article/8877278>

[Daneshyari.com](https://daneshyari.com)