

## Accepted Manuscript

Study of a new biocontrol fungal agent for avocado white root rot

Isabel Arjona-Girona, Carlos José López-Herrera

PII: S1049-9644(17)30184-6

DOI: <http://dx.doi.org/10.1016/j.biocontrol.2017.08.018>

Reference: YBCON 3642

To appear in: *Biological Control*

Received Date: 24 April 2017

Revised Date: 25 August 2017

Accepted Date: 29 August 2017



Please cite this article as: Arjona-Girona, I., López-Herrera, C.J., Study of a new biocontrol fungal agent for avocado white root rot, *Biological Control* (2017), doi: <http://dx.doi.org/10.1016/j.biocontrol.2017.08.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## STUDY OF A NEW BIOCONTROL FUNGAL AGENT FOR AVOCADO WHITE ROOT ROT

Isabel Arjona-Girona<sup>1</sup> and Carlos José López-Herrera\*

Departamento de Protección de Cultivos, Instituto de Agricultura Sostenible, C.S.I.C.  
C/Alameda del Obispo s/n, 14004, Córdoba, Spain.

\*Corresponding author: [lherrera@ias.csic.es](mailto:lherrera@ias.csic.es)

<sup>1</sup>[isabelarjona@ias.csic.es](mailto:isabelarjona@ias.csic.es)

### Abstract

Avocado white root rot is caused primarily by *Rosellinia necatrix* Prill. In this study, we buried bait twigs around avocado escape trees to recover non-pathogenic soil-borne fungi. First, we selected forty isolates: 19 were identified as *R. necatrix* and 21 as *Entoleuca* sp. The next step was to conduct pathogenicity tests on lupin and avocado plants, which showed that *R. necatrix* isolates were pathogenic in both types of plants, while *Entoleuca* sp. isolates were not. The optimal growth temperature *in vitro* for the *Entoleuca* sp. isolates was 30°C. *Dianthus caryophyllus* was found to be a host to this fungus, while *Olea europaea*, *Lupinus luteus*, *Asparagus officinalis* and *Fragaria ananassa* were not. Next, *Entoleuca* sp. was artificially inoculated into avocado plants and recovered from the roots up to two years later. In this way, we tested the biocontrol of the disease using *Entoleuca* sp. in avocado plants artificially inoculated with *R. necatrix*. Most of *Entoleuca* isolates (86%) controlled the disease, and we therefore concluded that they were effective biocontrol agents.

Download English Version:

<https://daneshyari.com/en/article/8877737>

Download Persian Version:

<https://daneshyari.com/article/8877737>

[Daneshyari.com](https://daneshyari.com)