



Impact of pest management practices on the frequency of insecticide resistance alleles in *Bemisia tabaci* (Hemiptera: Aleyrodidae) populations in three countries of West Africa

Olivier Gnankiné^{a,*}, Omer Hema^b, Moussa Namountougou^c, Laurence Mouton^d, Fabrice Vavre^d

^a Laboratoire d'Entomologie Appliquée, Université de Ouaga I Pr Joseph KI-ZERBO, Burkina Faso

^b Institut de l'Environnement et de Recherches Agricoles, BP 390 Bobo-Dioulasso, Burkina Faso

^c Université Nazi Boni de Bobo-Dioulasso, Burkina Faso

^d Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR5558, F-69622 Villeurbanne, France

ARTICLE INFO

Keywords:

Pest management programs

Insecticides

kdr

Ace1^R

Bemisia tabaci

ABSTRACT

In West Africa, the use of organophosphates and pyrethroid insecticides to control cotton pests has led to the evolution of resistance in field populations of *Bemisia tabaci* Gennadius. Three pest management programs have been commonly recommended: the Conventional Program (CP) where 6 treatments are applied, the use of *Bt* cotton plants for which only 2 applications of neonicotinoids are required and that has been adopted in many countries, and a biological program (BP) without any chemical treatment. The present study aimed to determine the influence of these practices on the frequency of mutations that confer resistance to pyrethroids (mutation L925I in the *para*-type voltage-gated sodium channel gene) and organophosphates (mutation F331W in the acetylcholinesterase enzyme *ace1*: allele *Ace1^R*) in *B. tabaci* populations using *Bt* cotton and CP areas in Pô and Saria (Burkina Faso), CP and BP areas in Kandi (Benin) and only CP areas in Tové and Infa (Togo). All individuals sampled belonged to the MED (biotypes MED-Q1) and Africa Silver Leafing (ASL) species. MED-Q1 was found in sympatry with ASL in Burkina Faso both on CP and *Bt* cotton areas at variable frequencies. In Togo and Benin, only ASL was found, except in Tové where MED-Q1 was also detected, but at low frequency. Frequencies of mutations that confer resistance varied between localities and species but we did not find any strong evidence of a relationship between the pest management program and these frequencies except for the allele *Ace1^R* in Burkina Faso for which the frequencies decrease when chemical applications are reduced. This study provides valuable information for the development of efficient integrated pest management programs.

1. Introduction

In western Africa, cotton is an economically important crop providing substantial incomes for farmers. However, the cotton plant is attacked by key pests including the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae) and the whitefly *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae). Three main spraying programs are recommended to control these pests in many countries of West Africa. They include the Conventional Cotton program (CP), the Biological Program (BP) and the transgenic *Bt* cotton program (*Bt* Cotton). The CP is based on calendar-based applications of insecticides belonging to pyrethroids, organophosphates and neonicotinoids families separately or as a mixture (Gnankiné et al., 2013a; Silvie et al., 2013). They are applied with temporal rotations during the whole

cotton season (from May to October). The repeated use of such insecticides have imposed strong selection pressures on target pests' populations, resulting in the evolution of field resistances (Houndété et al., 2010). Over \$60 millions of chemicals were spent for chemical pests control in Burkina Faso (Greenplate et al., 2003). The BP relies on the use of biopesticides and natural fertilizers without utilization of any chemical. The *Bt* cotton program uses *Bt* (*Bacillus thuringiensis*) transgenic cotton plants that express two crystal toxins (Cry1Ac and Cry2Ab) that target some major lepidopteran pests but are harmless to vertebrates and most other organisms (Mendelsohn et al., 2003; Sanahuja et al., 2011; Pardo-Lopez et al., 2013). In this program, only neonicotinoids are used at the end of the cotton phenological stages (Hémar et al., 2009). It was initiated in 2008 in West Africa but only in Burkina Faso. In 2016, the Burkina Faso government suspended the use of *Bt*

* Corresponding author. Laboratoire d'Entomologie Fondamentale et Appliquée, Département de Biologie et Physiologie Animales, Université de Ouagadougou, 03 BP 7021 Ouagadougou 03, Burkina Faso.

E-mail addresses: olivier.gnankine@univ-ouaga.bf, olgnankine@hotmail.com (O. Gnankiné).

<https://doi.org/10.1016/j.cropro.2017.10.020>

Received 20 September 2017; Received in revised form 30 October 2017; Accepted 31 October 2017

0261-2194/ © 2017 Elsevier Ltd. All rights reserved.

technology due to the quality of cotton fiber which is shorter than the fiber of conventional cotton.

One of major threat to cotton plant remains the whitefly *B. tabaci*. It causes damages directly through phloem feeding and indirectly through the transmission of plant viruses. *B. tabaci* is a complex of cryptic species whose taxonomy is still not entirely resolved. Based on the current consensus, *B. tabaci* is mostly represented by the MED species in western Africa, even though AnSL species can also be found (Gnankiné et al., 2013b, 2013c). Within the MED species, different biotypes are encountered including MED-Q1, MED-Q3 and ASL. Actually, recent analyses suggest that MED-Q1 and ASL do not hybridize in the field and that ASL is thus a different species (Mouton et al., 2015). Interestingly, these species/biotypes differ in terms of host plants range and insecticide resistance traits. Generally, MED-Q1 is predominant in cotton areas and sometimes found in sympatry with ASL on vegetables crops (Gnankiné et al., 2013b). It is also associated with higher levels of resistance to some insecticides as pyrethroids, organophosphates and neonicotinoids (Gnankiné et al., 2013a). In *B. tabaci*, two mutations in the *para*-type voltage-gated sodium channel gene, L925I and T929V, and one mutation in the acetylcholinesterase enzyme *ace1* (F331W: allele *Ace1^R*) confer resistance to pyrethroids and organophosphates, respectively (Roditakis et al., 2006; Alon et al., 2008; Tsagkarakou et al., 2009). Recent studies showed that the *Ace1^R* was found in both MED-Q1 and ASL but, while this resistant allele was almost fixed in MED-Q1 (0.99), its frequency was 0.59 in ASL (Mouton et al., 2015). In addition, while the L925I mutation in the sodium channel gene is almost fixed in MED-Q1 populations, it is rarely detected in ASL. The T929V was never found in *B. tabaci* populations from West Africa (Mouton et al., 2015). The objectives of the present study were to perform a first analysis of the impact of the agricultural practices on the *B. tabaci* biotypes/species composition and diversity, and the frequencies of alleles that confer resistance to pyrethroids and organophosphates. Sampling was performed in three countries of Western Africa: Burkina Faso, Benin and Togo.

2. Materials and methods

2.1. Management of cotton pests

Three pest control programs are recommended in West African countries:

- The Conventional Program (CP) is based on two to four treatments with pyrethroids (PY) plus organophosphates (OP) and 2 other treatments with neonicotinoids (see Table 1 for details).
- The Biological control Program (BP) does not use chemicals for plant protection. Farmers worked under the supervision of technicians from the Beninese Organization for Organic Farming Promotion (OBEPAP) who participated in the implementation and the survey of good agricultural practices on organic cotton.
- The transgenic cotton program (*Bt* Cotton). In this program, pesticides belonging to OP and PYR are not used. Only neonicotinoids are used at boll opening stage to control sucking pests. In this case, farmers worked under the supervision of technicians from societies of textile fibers in Burkina Faso.

2.2. *B. tabaci* sampling

Sampling was performed in october and november between 2009 and 2015 in three countries of Western Africa: Burkina Faso, Benin and Togo (Fig. 1, Table 2). In Burkina Faso, whiteflies were collected randomly in two localities, Pô in 2013 and Saria in 2015, from *Bt* cotton and CP fields. In Pô, *Bt* cotton represented 90% of the sampled fields while in Saria it represented 15%. In Benin, sampling was done at Kandi in 2009 in CP and BP areas (BP represent around 10% of areas). In Togo, collection was done randomly in two sites in 2009, Infa and Tové,

where only CP is used. The collected adult whiteflies were stored in ethanol 95%. The origin of the samples (location) and the number of individuals are summarized in Table 2.

2.3. Molecular analysis

2.3.1. DNA extraction

For each individual, total DNA was extracted in 25 µl of an extraction buffer containing 50 mM KCl, 10 mM Tris-base pH 8, 0.45% Nonidet P-40, 0.45% Tween 20 and 50 mg/ml proteinase K. After 3 h at 65 °C, samples were incubated at 100 °C for 15 min. Pure water (35 µl) was then added to the extract.

2.3.2. Identification of *B. tabaci*

Species/biotypes were identified using the Polymerase Chain Reaction-Random Fragment Length Polymorphism (PCR-RFLP) diagnostic assay based on the mitochondrial cytochrome oxidase 1 gene sequence (*mtCO1*) described in Henri et al. (2013). This technique allows discriminating the species/biotypes present in West Africa (Gnankiné et al., 2013b).

2.3.3. Identification of susceptible and resistant alleles of the sodium channel and the acetylcholinesterase *ace1* genes

Resistant and susceptible alleles in the *para*-type voltage-gated sodium channel and *ace1* genes were identified using the diagnostic assays developed by Tsagkarakou et al. (2009). Briefly, *ace1*-susceptible (F331) and -resistant (W331) alleles, as well as susceptible (L925) and resistant (I925) *para*-type voltage-gated sodium channel alleles were detected using PCR-RFLP (Tsagkarakou et al., 2009). Some PCR products were sequenced for each susceptible and resistant allele and each country. We never found the T929V mutation in the sequences.

The frequencies of *kdr* and *ace-1^R* mutations were calculated according to the formula.

$$p = \frac{n\sigma^{\circ}(R) + 2n\sigma^{\circ}(RR) + n\sigma^{\circ}(RS)}{n\sigma^{\circ} + 2n\sigma^{\circ}} \text{ where } RR \text{ was the number of homozygotes, } RS \text{ the number of heterozygotes and } n \text{ the size of specimens analysed.}$$

2.4. Statistical analyses

Statistical analysis were performed using the R statistical software (<http://www.R-project.org>). The effects of the pest management practices on the proportions of *B. tabaci* composition and the frequencies of resistance alleles were tested by using Fisher's exact tests.

3. Results

3.1. Geographic distribution of biotypes

All the 170 *B. tabaci* individuals collected in 5 localities in Burkina Faso, Benin and Togo belonged to MED-Q1 or ASL (Table 2). In Togo and Benin, only ASL was found (except one MED-Q1 individual in Tové), while in Burkina Faso, MED-Q1 and ASL were found in sympatry at variable frequencies: depending on the locality, it was either ASL or MED-Q1 that predominated. In Pô, the frequency of ASL reached more than 80% whatever the control strategy (CP or Cotton *Bt*). In Saria, MED-Q1 was more common than ASL, but their relative frequencies depended on the management program: on CP areas, 93% of individuals belonged to MED-Q1 while only 57% of whiteflies were MED-Q1 on *Bt* Cotton fields (Fisher exact test, $p < 0.005$).

3.2. Frequency of the L925I mutation

For the *para*-type voltage-gated sodium channel gene, we studied the frequency of the L925I mutation that correspond to the allele called r1 by Alon et al. (2008). We found high variations depending on the country (Fisher exact test, $p < 0.0005$). Indeed, in Burkina Faso, r1

Download English Version:

<https://daneshyari.com/en/article/8878327>

Download Persian Version:

<https://daneshyari.com/article/8878327>

[Daneshyari.com](https://daneshyari.com)