

## Accepted Manuscript

Global estimation of long-term persistence in annual river runoff

Y. Markonis , Y. Moustakis , C. Nasika , P. Sychova ,  
P. Dimitriadis , M. Hanel , P. Máca , S.M. Papalexiou

PII: S0309-1708(17)30364-0  
DOI: [10.1016/j.advwatres.2018.01.003](https://doi.org/10.1016/j.advwatres.2018.01.003)  
Reference: ADWR 3059



To appear in: *Advances in Water Resources*

Received date: 11 April 2017  
Revised date: 2 January 2018  
Accepted date: 3 January 2018

Please cite this article as: Y. Markonis , Y. Moustakis , C. Nasika , P. Sychova , P. Dimitriadis , M. Hanel , P. Máca , S.M. Papalexiou , Global estimation of long-term persistence in annual river runoff, *Advances in Water Resources* (2018), doi: [10.1016/j.advwatres.2018.01.003](https://doi.org/10.1016/j.advwatres.2018.01.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights:**

- Hurst coefficient  $H = 0.65$ , as estimated in 696 annual, globally distributed, streamflow records with at least 80 years of data.
- Long-term persistence in annual streamflow is weaker than previous estimates.
- The estimated values of  $H$  can be reproduced by an auto-regressive AR(1) process.
- Some of the most common methods for the estimation of  $H$  coefficient, might not be able to discriminate short- and long-term persistence.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8883343>

Download Persian Version:

<https://daneshyari.com/article/8883343>

[Daneshyari.com](https://daneshyari.com)