


Author's Accepted Manuscript

Bottom-trawling fisheries influence on standing stocks, composition, diversity and trophic redundancy of macrofaunal assemblages from the West Iberian Margin

Sofia Pinto Ramalho, Mariana Almeida, Patricia Esquete, Luciana Génio, Ascensão Ravara, Clara F. Rodrigues, Nikolaos Lampadariou, Ann Vanreusel, Marina R. Cunha



www.elsevier.com

PII: S0967-0637(18)30001-3
DOI: <https://doi.org/10.1016/j.dsr.2018.06.004>
Reference: DSR12921

To appear in: *Deep-Sea Research Part I*

Received date: 5 January 2018
Revised date: 16 May 2018
Accepted date: 8 June 2018

Cite this article as: Sofia Pinto Ramalho, Mariana Almeida, Patricia Esquete, Luciana Génio, Ascensão Ravara, Clara F. Rodrigues, Nikolaos Lampadariou, Ann Vanreusel and Marina R. Cunha, Bottom-trawling fisheries influence on standing stocks, composition, diversity and trophic redundancy of macrofaunal assemblages from the West Iberian Margin, *Deep-Sea Research Part I*, <https://doi.org/10.1016/j.dsr.2018.06.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Bottom-trawling fisheries influence on standing stocks, composition, diversity and trophic redundancy of macrofaunal assemblages from the West Iberian Margin

Sofia Pinto Ramalho^{1,2*}, Mariana Almeida¹, Patricia Esquete¹, Luciana Génio¹, Ascensão Ravara¹, Clara F. Rodrigues¹, Nikolaos Lampadariou³, Ann Vanreusel², Marina R. Cunha¹

¹Departamento de Biologia, Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

²Department of Biology, Marine Biology Research Group, Ghent University, Krijgslaan 281/S8, 9000 Ghent, Belgium

³Hellenic Centre for Marine Research, Institute of Oceanography, P.O. Box 2214, 71003, Heraklion, Crete, Greece

Email address: sofia.pinto.ramalho@gmail.com;
sramalho@ua.pt

*Corresponding author: Sofia P. Ramalho

Abstract

Bottom-trawling fisheries operating in Portugal (West Iberian Margin) impose one of the largest footprints per unit of biomass landed in European waters at depths greater than 200 m, affecting the seafloor integrity and the associated benthic fauna. To investigate how trawling pressure is affecting the macrofaunal assemblages, we compared the standing stock (abundance and biomass), community structure and taxonomical and trophic diversity in areas subjected to varying trawling pressure along the SW Portuguese upper slope, between 200-600 m. In addition to trawling pressure, several environmental variables, namely depth, grain size and organic matter, were correlated with the biological component, which suggest that the longstanding trawling pressure presents cumulative effects to the habitat heterogeneity known to characterise the West Iberian Margin fauna. Furthermore, our results showed a depletion of macro-infaunal abundances in both the fishing ground and the adjacent area (up to 3 times lower), when compared to the area not trawled. The observed decrease in abundance with increasing trawling pressure was also associated with a loss of species and trophic richness, but univariate diversity indices

Download English Version:

<https://daneshyari.com/en/article/8884206>

Download Persian Version:

<https://daneshyari.com/article/8884206>

[Daneshyari.com](https://daneshyari.com)