Accepted Manuscript

Chlorogenic and phenolic acids are only very weak inhibitors of human salivary α -amylase and rat intestinal maltase activities

Hilda Nyambe-Silavwe, Gary Williamson



PII:	80963-9969(18)30576-3
DOI:	doi:10.1016/j.foodres.2018.07.038
Reference:	FRIN 7778
To appear in:	Food Research International
Received date:	25 May 2018
Revised date:	27 June 2018
Accepted date:	28 July 2018

Please cite this article as: Hilda Nyambe-Silavwe, Gary Williamson , Chlorogenic and phenolic acids are only very weak inhibitors of human salivary α -amylase and rat intestinal maltase activities. Frin (2018), doi:10.1016/j.foodres.2018.07.038

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Chlorogenic and phenolic acids are only very weak inhibitors of human

salivary α -amylase and rat intestinal maltase activities

Hilda Nyambe-Silavwe and Gary Williamson

School of Food Science and Nutrition, University of Leeds, Leeds, LS2 9JT, UK

Correspondence to G. Williamson (g.williamson@leeds.ac.uk)

Abbreviations:

5-CQA: 5-caffeoylquinic acid

PBS: phosphate-buffered saline

DNS: 3,5-dinitrosalicylic acid

Download English Version:

https://daneshyari.com/en/article/8888534

Download Persian Version:

https://daneshyari.com/article/8888534

Daneshyari.com