

## Accepted Manuscript

Reprint of: Boundary conditions for fractional diffusion

Boris Baeumer, Mihály Kovács, Mark M. Meerschaert,  
Harish Sankaranarayanan



PII: S0377-0427(18)30136-5  
DOI: <https://doi.org/10.1016/j.cam.2018.03.007>  
Reference: CAM 11557

To appear in: *Journal of Computational and Applied  
Mathematics*

Received date : 26 May 2017  
Revised date : 18 December 2017

Please cite this article as: B. Baeumer, M. Kovács, M.M. Meerschaert, H. Sankaranarayanan,  
Reprint of: Boundary conditions for fractional diffusion, *Journal of Computational and Applied  
Mathematics* (2018), <https://doi.org/10.1016/j.cam.2018.03.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## BOUNDARY CONDITIONS FOR FRACTIONAL DIFFUSION

BORIS BAEUMER, MIHÁLY KOVÁCS, MARK M. MEERSCHAERT,  
AND HARISH SANKARANARAYANAN

ABSTRACT. This paper derives physically meaningful boundary conditions for fractional diffusion equations, using a mass balance approach. Numerical solutions are presented, and theoretical properties are reviewed, including well-posedness and steady state solutions. Absorbing and reflecting boundary conditions are considered, and illustrated through several examples. Reflecting boundary conditions involve fractional derivatives. The Caputo fractional derivative is shown to be unsuitable for modeling fractional diffusion, since the resulting boundary value problem is not positivity preserving.

## 1. INTRODUCTION

The space-fractional diffusion equation replaces the second derivative or Laplacian in the traditional diffusion equation with a fractional derivative. Fractional derivatives were invented soon after their integer-order counterparts, and by now have become an established field of study with a wide variety of applications in science and technology [26, 34, 45, 44, 53, 54, 61, 65]. Practical applications include physics [27, 41, 60, 72], finance [23, 32, 46, 66, 67], microbiology [6, 20, 29], medical imaging [8, 42, 43, 58, 73, 84], and hydrology [10, 15, 33, 9]. Zaslavsky [81, 82] initiated the application of fractional calculus to chaotic dynamical systems. These methods may also be applicable to nonlinear dynamics in Hodgkin-Huxley neurons and pancreatic beta cells [24, 25, 69]. Many effective numerical methods have been developed for fractional differential equations, along with proofs of stability and consistency [7, 14, 16, 22, 39, 40, 47, 48, 49, 56, 61, 62, 71, 76, 80, 83]. Because fractional

---

*Date:* December 18, 2017.

*Key words and phrases.* Fractional calculus; boundary value problem; numerical solution; well-posed.

Baeumer was partially supported by the Marsden Fund administered by the Royal Society of New Zealand.

Kovács was partially supported by the Marsden Fund administered by the Royal Society of New Zealand.

Meerschaert was partially supported by ARO MURI grant W911NF-15-1-0562 and NSF grants DMS-1462156 and EAR-1344280.

Sankaranarayanan was supported by ARO MURI grant W911NF-15-1-0562.

Download English Version:

<https://daneshyari.com/en/article/8902052>

Download Persian Version:

<https://daneshyari.com/article/8902052>

[Daneshyari.com](https://daneshyari.com)