ARTICLE IN PRESS

C. R. Acad. Sci. Paris, Ser. I ••• (••••) •••-•••



Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I



CRASS1:5998

www.sciencedirect.com

Mathematical analysis/Mathematical physics

Existence and uniqueness of solutions to a model describing miscible liquids

Existence et unicité des solutions pour un modèle décrivant les liquides miscibles

Karam Allali

Laboratory Mathematics and Applications, Department of Mathematics, Faculty of Sciences and Technologies, University Hassan II of Casablanca, PO Box 146, Mohammedia, Morocco

ARTICLE INFO

Article history: Received 20 August 2017 Accepted after revision 18 October 2017 Available online xxxx

Presented by the Editorial Board

ABSTRACT

The existence and the uniqueness of solutions to a problem of miscible liquids are investigated in this note. The model consists of Navier–Stokes equations with Korteweg stress terms coupled with the reaction–diffusion equation for the concentration. We assume that the fluid is incompressible and the Boussinesq approximation is adopted. The global existence and uniqueness of solutions is established for some optimal conditions on the reaction source term and the external force functions.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

RÉSUMÉ

Dans cette note, nous étudierons un problème d'existence et d'unicité pour un modèle qui décrit l'interaction de deux fluides miscibles. Le modèle considéré prend la forme d'équations de Navier–Stokes avec contraintes de Korteweg couplées à l'équation de réaction– diffusion de la concentration. Nous supposons que le fluide est incompressible; l'approximation de Boussinesq est adoptée. L'existence globale et l'unicité des solutions sont établies pour des conditions optimales sur le terme source de réaction et les forces externes. © 2017 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

The theoretical and experimental investigation of the dynamics of two miscible liquids have attracted considerable attention from many researchers because of its significant applications in various fields such as enhanced oil recovery, hydrology, frontal polymerization, groundwater pollution, and filtration [1,2,6]. A detailed review of the results on the interfacial dy-

https://doi.org/10.1016/j.crma.2017.10.009

1631-073X/ $\! \textcircled{\odot}$ 2017 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

E-mail address: allali@fstm.ac.ma.

00

2

ARTICLE IN PRESS

K. Allali / C. R. Acad. Sci. Paris, Ser. I ••• (••••) •••-•••

namics of two miscible liquids is presented in [3]. In this paper, we are interested in studying the following miscible liquids model:

$$\frac{\partial C}{\partial t} + u \cdot \nabla C = d\Delta C - Cg,\tag{1.1}$$

$$\frac{\partial u}{\partial t} + (u \cdot \nabla)u = -\nabla p + \mu \Delta u + \nabla \cdot T(C) + f,$$
(1.2)

$$\operatorname{div}(u) = 0. \tag{1.3}$$

with the following boundary conditions:

$$\frac{\partial C}{\partial n} = 0, \ u = 0 \ \text{on } \Gamma,$$
 (1.4)

and the following initial conditions:

$$C(x,0) = C_0(x), \ u(x,0) = u_0(x), \ x \in \Omega.$$
(1.5)

Here *u* is the velocity, *p* is the pressure, *C* is the concentration, *d* is the coefficient of mass diffusion, μ is the viscosity, Γ is a Lipschitz continuous boundary of the open bounded domain Ω , *n* is the unit outward normal vector to Γ , *f* is the function describing the external forces such as gravity and buoyancy, while *g* stands for the source term. The Korteweg stress tensor terms are given by the relations [4]:

$$T_{11} = k \left(\frac{\partial C}{\partial x_2}\right)^2, \ T_{12} = T_{21} = -k \frac{\partial C}{\partial x_1} \frac{\partial C}{\partial x_2}, \ T_{13} = T_{31} = -k \frac{\partial C}{\partial x_1} \frac{\partial C}{\partial x_3},$$
$$T_{23} = T_{32} = -k \frac{\partial C}{\partial x_2} \frac{\partial C}{\partial x_3}, \ T_{22} = k \left(\frac{\partial C}{\partial x_1}\right)^2, \ T_{33} = k \left(\frac{\partial C}{\partial x_3}\right)^2,$$

here k is a nonnegative constant. We set

$$\nabla \cdot T(C) = \begin{pmatrix} \frac{\partial T_{11}}{\partial x_1} + \frac{\partial T_{12}}{\partial x_2} + \frac{\partial T_{13}}{\partial x_3} \\ \frac{\partial T_{21}}{\partial x_1} + \frac{\partial T_{22}}{\partial x_2} + \frac{\partial T_{23}}{\partial x_3} \\ \frac{\partial T_{31}}{\partial x_1} + \frac{\partial T_{32}}{\partial x_2} + \frac{\partial T_{33}}{\partial x_3} \end{pmatrix}.$$

In the absence of external and source forces (f = g = 0) and in the two-dimensional case, the existence and uniqueness of a solution to the model (1.1)–(1.5) is studied in [5]. The aim of this paper is to continue the investigations of miscible liquids by considering three-dimensional Navier–Stokes equations and by introducing the external forces in the equation of motion and the source terms in the equation for the concentration.

2. Existence of solutions

First, we will introduce our functional framework on which the study will be carried out. Let the following velocity and concentration spaces:

$$S_u = \{u \in H_0^1(\Omega); \operatorname{div}(u) = 0\}, \quad S_C = \{C \in H^2(\Omega); \frac{\partial C}{\partial n} = 0 \text{ on } \Gamma\}.$$

Hence, the variational formulation of the problem is to find $C \in S_C$ and also $u \in S_u$ such that, for all $B \in S_C$ and $v \in S_u$, we have the following equalities:

$$\left(\frac{\partial C}{\partial t}, B\right) + d(\nabla C, \nabla B) + (u \cdot \nabla C, B) + (gC, B) = 0,$$

$$\left(\frac{\partial u}{\partial t}, v\right) + \mu(\nabla u, \nabla v) - (\operatorname{div} T(C), v) - (f, v) = 0.$$
(2.2)

In what follows, we will assume that the function f(x, t) is positive and bounded in $L^{\infty}(0, t; L^2(\Omega))$; we will assume also that g(x) is positive and bounded in $L^{\infty}(\Omega)$. In other words, there exist two real positive constants \overline{f} and \overline{g} , such that

$$\|f\|_{L^{\infty}(0,t;L^{2}(\Omega))} \le f, \ f(x,t) \ge 0$$
(2.3)

and

$$\|g\|_{L^{\infty}(\Omega)} \leq \overline{g}, \quad g(x) \geq 0.$$

$$(2.4)$$

In order to prove the existence of solutions, we will need the following lemmas.

Please cite this article in press as: K. Allali, Existence and uniqueness of solutions to a model describing miscible liquids, C. R. Acad. Sci. Paris, Ser. I (2017), https://doi.org/10.1016/j.crma.2017.10.009

Download English Version:

https://daneshyari.com/en/article/8905597

Download Persian Version:

https://daneshyari.com/article/8905597

Daneshyari.com