

Hypoxia in the Holocene Baltic Sea: Comparing modern versus past intervals using sedimentary trace metals

Niels A.G.M. van Helmond, Tom Jilbert, Caroline P. Slomp

PII: S0009-2541(18)30330-9
DOI: doi:10.1016/j.chemgeo.2018.06.028
Reference: CHEMGE 18827

To appear in: *Chemical Geology*

Received date: 13 April 2018
Revised date: 25 June 2018
Accepted date: 29 June 2018

Please cite this article as: Niels A.G.M. van Helmond, Tom Jilbert, Caroline P. Slomp , Hypoxia in the Holocene Baltic Sea: Comparing modern versus past intervals using sedimentary trace metals. Chemge (2018), doi:10.1016/j.chemgeo.2018.06.028

Hypoxia in the Holocene Baltic Sea: comparing modern versus past intervals using sedimentary trace metals

Niels A.G.M. van Helmond[a*], Tom Jilbert[a,b] and Caroline P. Slomp[a]

*Corresponding author: E-mail: n.vanhelmond@uu.nl

[a] Department of Earth Sciences, Faculty of Geosciences, Utrecht University. Princetonlaan 8a, 3584 CB Utrecht, Netherlands; n.vanhelmond@uu.nl (Niels A.G.M. van Helmond), c.p.slomp@uu.nl (Caroline P. Slomp)

[b] Ecosystems and Environment Research Program, Faculty of Biological and Environmental Sciences, University of Helsinki, Finland; tom.jilbert@helsinki.fi