



Malaysia Airlines flight MH370 search data reveal geomorphology and seafloor processes in the remote southeast Indian Ocean

Kim Picard^{a,*}, Brendan P. Brooke^a, Peter T. Harris^b, Paulus J.W. Siwabessy^a, Millard F. Coffin^{c,d,e}, Maggie Tran^a, Michele Spinoccia^a, Jonathan Weales^a, Miles Macmillan-Lawler^b, Jonah Sullivan^a

^a Geoscience Australia, Environmental Geoscience Division, GPO Box 378, Canberra, ACT 2601, Australia

^b GRID-Arendal, Postboks 183, N-4802 Arendal, Norway

^c Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 129, Hobart, Tasmania 7001, Australia

^d School of Earth and Climate Sciences, University of Maine, Orono, ME 04473, USA

^e Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

ARTICLE INFO

Keywords:

Indian Ocean
Multibeam echosounder
Geomorphology
Processes
Deepsea
Seamount

ABSTRACT

A high-resolution multibeam echosounder (MBES) dataset covering over 279,000 km² was acquired in the southeastern Indian Ocean to assist the search for Malaysia Airlines Flight 370 (MH370) that disappeared on 8 March 2014. The data provided an essential geospatial framework for the search and is the first large-scale coverage of MBES data in this region. Here we report on geomorphic analyses of the new MBES data, including a comparison with the Global Seafloor Geomorphic Features Map (GSFM) that is based on coarser resolution satellite altimetry data, and the insights the new data provide into geological processes that have formed and are currently shaping this remote deepsea area. Our comparison between the new MBES bathymetric model and the latest global topographic/bathymetric model (SRTM15_plus) reveals that 62% of the satellite-derived data points for the study area are comparable with MBES measurements within the estimated vertical uncertainty of the SRTM15_plus model (± 100 m). However, > 38% of the SRTM15_plus depth estimates disagree with the MBES data by > 100 m, in places by up to 1900 m. The new MBES data show that abyssal plains and basins in the study area are significantly more rugged than their representation in the GSFM, with a 20% increase in the extent of hills and mountains. The new model also reveals four times more seamounts than presented in the GSFM, suggesting more of these features than previously estimated for the broader region. This is important considering the ecological significance of high-relief structures on the seabed, such as hosting high levels of biodiversity. Analyses of the new data also enabled sea knolls, fans, valleys, canyons, troughs, and holes to be identified, doubling the number of discrete features mapped.

Importantly, mapping the study area using MBES data improves our understanding of the geological evolution of the region and reveals a range of modern sedimentary processes. For example, a large series of ridges extending over approximately 20% of the mapped area, in places capped by sea knolls, highlight the preserved seafloor spreading fabric and provide valuable insights into Southeast Indian Ridge seafloor spreading processes, especially volcanism. Rifting is also recorded along the Broken Ridge – Diamantina Escarpment, with rift blocks and well-bedded sedimentary bedrock outcrops discernible down to 2400 m water depth. Modern ocean floor sedimentary processes are documented by sediment mass transport features, especially along the northern margin of Broken Ridge, and in pockmarks (the finest-scale features mapped), which are numerous south of Diamantina Trench and appear to record gas and/or fluid discharge from underlying marine sediments. The new MBES data highlight the complexity of the search area and serve to demonstrate how little we know about the vast areas of the ocean that have not been mapped with MBES. The availability of high-resolution and accurate maps of the ocean floor can clearly provide new insights into the Earth's geological evolution, modern ocean floor processes, and the location of sites that are likely to have relatively high biodiversity.

* Corresponding author.

E-mail address: kim.picard@ga.gov.au (K. Picard).

<https://doi.org/10.1016/j.margeo.2017.10.014>

Received 9 July 2017; Received in revised form 24 October 2017; Accepted 29 October 2017

Available online 08 November 2017

0025-3227/ Crown Copyright © 2017 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).




Fig. 1. Location map showing the multibeam bathymetry data combined with sun-illuminated relief collected in the search for MH370. The locations of Figs. 2 and 6 to 9 are displayed as well as the locations of DSDP and ODP drill sites 255 (Leg 26), 752–755 (Leg 121), and 1141/1142 (Leg 183). Upper left inset shows a map of the multibeam bathymetric coverage (pink) overlaid on the SRTM15_plus model (Olson et al., 2016). The map highlights the Southeast Indian Ridge (SEIR), estimated spreading rates of the SEIR (Argus et al., 2011), the magnetic isochrons (Müller et al., 2008), and interpreted SEIR segments (Small et al., 1999). SWIR: Southwest Indian Ridge; CIR: Central Indian Ridge; RTJ: Rodriguez Triple Junction; AAD: Australian–Antarctic Discordance; WA: Western Australia. Lower right inset is an excerpt of the Global Seafloor Geomorphic Feature Map (GSFM; Harris et al., 2014) corresponding to the search area. PP-715-1 is Geoscience Australia internal production number for this figure. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

1. Introduction

Malaysia Airlines Flight 370 (MH370) was a scheduled international passenger flight that disappeared on 8 March 2014 while flying from

Kuala Lumpur, Malaysia, to Beijing, China. In the ensuing multi-national effort to search for aircraft wreckage, a large area of ocean floor was mapped using ship-mounted multibeam echosounders (MBES), providing unprecedented detail in an otherwise remote area of the

Download English Version:

<https://daneshyari.com/en/article/8912044>

Download Persian Version:

<https://daneshyari.com/article/8912044>

[Daneshyari.com](https://daneshyari.com)