

Accepted Manuscript

Quaternary history of the Kiseiba Oasis region, southern Egypt

Ted A. Maxwell, C. Vance Haynes, Kathleen Nicoll, Andrew K. Johnston, John A. Grant, Ali Kilani



PII: S1464-343X(17)30168-1
DOI: 10.1016/j.jafrearsci.2017.04.024
Reference: AES 2892
To appear in: *Journal of African Earth Sciences*
Received Date: 31 October 2016
Revised Date: 02 April 2017
Accepted Date: 20 April 2017

Please cite this article as: Ted A. Maxwell, C. Vance Haynes, Kathleen Nicoll, Andrew K. Johnston, John A. Grant, Ali Kilani, Quaternary history of the Kiseiba Oasis region, southern Egypt, *Journal of African Earth Sciences* (2017), doi: 10.1016/j.jafrearsci.2017.04.024

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Fluvial channels detected by radar are correlated with near-surface sedimentary units

Inverted topography coupled with paleodrainage is responsible for the these channels

Quaternary drainage from south to north was responsible for fluvial deposits

Topographic inversion led to local drainage reversal in the late Pleistocene

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8913738>

Download Persian Version:

<https://daneshyari.com/article/8913738>

[Daneshyari.com](https://daneshyari.com)