Accepted Manuscript

Revision of the Pennsylvanian fern *Boweria* Kidston and the establishment of the new genus *Kidstoniopteris*

Jana Frojdová, Josef Pšenička, Jiří Bek, Christopher J. Cleal

PII: S0034-6667(15)30005-1

DOI: doi:10.1016/j.revpalbo.2016.08.011

Reference: PALBO 3787

To appear in: Review of Palaeobotany and Palynology

Received date: 12 August 2015 Revised date: 13 July 2016 Accepted date: 31 August 2016



Please cite this article as: Frojdová, Jana, Pšenička, Josef, Bek, Jiří, Cleal, Christopher J., Revision of the Pennsylvanian fern *Boweria* Kidston and the establishment of the new genus *Kidstoniopteris*, *Review of Palaeobotany and Palynology* (2016), doi:10.1016/j.revpalbo.2016.08.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Revision of the Pennsylvanian fern *Boweria* Kidston and the establishment of the new genus *Kidstoniopteris*

Jana Frojdová ^{a, b}*, Josef Pšenička ^c, Jiří Bek ^a, Christopher J. Cleal ^d

^a Laboratory of Palaeobiology and Palaeoecology, Institute of Geology of the Czech Academy of Sciences, v.v.i., Rozvojová 269,165 00 Prague 6, Czech Republic

^b Institute of Geology and Palaeontology, Charles University in Prague, Faculty of Science, Albertov 6, 128 43 Prague 2, Czech Republic

^c Centre of Palaeobiodiversity, West Bohemia Museum in Pilsen, Kopeckého sady 2, Pilsen 301 00, Czech Republic

^d Department of Natural Sciences, National Museum Wales, Cardiff, CF10 3NP, United Kingdom

*corresponding author: frojdova@gli.cas.cz

Abstract

Boweria is a fossil-genus of Pennsylvanian true ferns whose exact taxonomic position was for many years confused. The morphology of reproductive structures (including sporangia), pinnae, pinnules and in situ spores (Leiotriletes and Granulatisporites types) of the type species (Boweria schatzlarensis) are described in detail. A new species (Boweria nowarudensis) is distinguished by its sporangia being isolated with two types of

Download English Version:

https://daneshyari.com/en/article/8916658

Download Persian Version:

https://daneshyari.com/article/8916658

<u>Daneshyari.com</u>