

# Clot or Not?



Maude Peretz-Larochelle, MD, Sarah Blissett, MD, MHPE, Jed Lipes, MD, Bojan Kovacina, MD, Jonathan Afilalo, MD, MSc, and Lawrence Rudski, MD, *Montreal, Quebec, and London, Ontario, Canada*

## INTRODUCTION

Cardiac masses are rare, with a prevalence of 0.15% in echocardiographic series.<sup>1</sup> The differential diagnosis for cardiac masses includes primary cardiac tumors, cardiac metastases, vegetations, thrombi, fluid-filled lesions, and artifacts.<sup>1</sup> These entities have widely different implications for management and prognosis, highlighting the importance of accurately identifying the etiology of a cardiac mass. Echocardiography is a key tool for the identification and diagnosis of cardiac masses, relying on the appearance and location of the mass. Multimodality imaging permits the integration of echocardiography and other noninvasive imaging modalities to improve the characterization and differentiation of cardiac masses. This integrated approach allows tissue characterization without the use of a cardiac biopsy and its associated risks.

We report a case of multiple right ventricular (RV) cardiac masses characterized using an integrated imaging approach.

## CASE PRESENTATION

An 85-year-old cachectic man, an ex-smoker, presented to the emergency department with a 3-month history of progressive exertional dyspnea. His medical history was notable for laryngeal cancer resected in 2010 and recent diagnosis of invasive urothelial carcinoma without prostatic invasion, status post resection with curative intent 3 months before presentation.

His blood pressure on arrival was 121/64 mm Hg, with a heart rate of 77 beats/min and mild hypoxemia. Laboratory values revealed stable high-sensitivity troponin T of 20 ng/L (upper limit of normal, 15 ng/L) and prostate-specific antigen of 1.2  $\mu\text{g/L}$  (upper limit of normal, 4.0  $\mu\text{g/L}$ ).

On point-of-care ultrasound, the emergency physician identified RV masses, raising the differential diagnosis of tumors versus thrombi. Computed tomographic angiography described extensive bilateral lobar and segmental arterial filling defects compatible with pulmonary emboli, accompanied by masslike filling defects

within the RV cavity and features of right heart strain. Low-dose thrombolysis was considered but withheld because of the non-thrombotic echocardiographic appearance of the masses and the patient's hemodynamic stability.

Formal transthoracic echocardiography revealed moderate RV enlargement with global hypokinesis and multiple large masses in all segments of the right ventricle (inflow, body, and outflow). Masses were adherent to the lateral free wall, the septal aspect of the tricuspid annulus, the papillary muscles, and the RV apex (Figures 1 and 2, Videos 1 and 2).

The masses were multilobulated and had similar echogenicity to the myocardium without independent mobility. No right atrial or inferior vena cava masses were observed (Figure 3).

There was secondary mild tricuspid valve inflow obstruction (mean transtricuspid gradient, 2 mm Hg at 80 beats/min) (Figure 4) and moderate tricuspid regurgitation (Video 3), with an estimated pulmonary artery systolic pressure of 59 mm Hg. There was a small circumferential pericardial effusion without echocardiographic features of tamponade (Figure 1, Video 1).

The constellation of these echocardiographic findings suggested tumor rather than multiple thrombi; however, superimposed thrombi could not be excluded.

To further characterize the masses, cardiac magnetic resonance imaging (MRI) was performed. Cardiac MRI confirmed the location and burden of the RV masses (Figures 5 and 6, Videos 4 and 5).

Tissue characteristics were consistent with tumor without superimposed thrombi: T1 isointense (Figure 7A), T2 hyperintense (Figure 7B), first-pass perfusion positive (Figure 7C), and late gadolinium enhancement (LGE) heterogeneously positive.

The patient died of his disease within 1 month of diagnosis of cardiac involvement.

## DISCUSSION

Echocardiography is the first-line imaging modality for cardiac masses because it is a portable, widely available, noncontrast, and dynamic technology. It can also be used by nonechocardiographers to enhance their clinical assessments.<sup>2</sup>

In this case, in view of the point-of-care ultrasound and computed tomographic results, a presumptive diagnosis of multiple pulmonary emboli was made, and the emergency department physician considered low-dose thrombolysis<sup>3</sup> on the basis of the "clot" burden and RV strain. Appropriately, formal echocardiography was performed given the uncertainty of the etiology of the masses. The imaging findings on echocardiography, particularly the heterogeneity of the RV masses, their distribution, and motion synchronous with myocardium, suspected invasion of the RV lateral wall, as well as the thickened pericardium, suggested tumor rather than thrombi. Therefore, thrombolysis was deferred on the basis of the echocardiographic findings.

Considering our patient's history of recent urothelial carcinoma, cardiac metastases were strongly suspected. We proceeded to

From the Division of Cardiology (M.P.-L., S.B., J.A., L.R.), the Department of Critical Care (J.L.), and the Department of Radiology (B.K.), Jewish General Hospital, Montreal, Quebec, Canada; and the Division of Cardiology, London Health Science Centre, London, Ontario, Canada (S.B.).


Keywords: Right ventricular masses, Cardiac tumor, Cardiac thrombus, Cardiac metastases

Conflicts of interest: Dr. Rudski has a minor holding in GE stock outside of a managed portfolio. All other authors reported no actual or potential conflicts of interest relative to this document.


Copyright 2017 by the American Society of Echocardiography. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2468-6441

<https://doi.org/10.1016/j.case.2017.09.010>




**Figure 1** Apical four-chamber view demonstrating a heterogeneous mass adjacent to the lateral RV free wall, possibly invading the myocardium (*star*). There were separate heterogeneous, multilobed masses on the papillary muscles (*asterisk*). Moderate RV dilation was noted. There was a small pericardial effusion.




**Figure 4** Mild RV inflow obstruction (mean transtricuspid gradient, 2 mm Hg at 80 beats/min).




**Figure 2** Off-axis view demonstrating the masses on the papillary muscles in the RV body (*asterisk*) and a separate mass originating in the RV apex extending into the right ventricular outflow tract (*solid circle*). The masses were not independently mobile.



**Figure 3** Plethoric inferior vena cava (25 mm) without luminal masses.



**Figure 5** RV inflow and outflow steady-state free precession MRI sequence demonstrating the RV masses isointense to the myocardium originating from the base of the papillary muscles (*asterisk*), right ventricular outflow tract (*solid circle*), RV lateral wall (*star*), and tricuspid valve annulus (*diamond*).



**Figure 6** Four-chamber steady-state free precession MRI sequence demonstrating the RV masses isointense to the myocardium originating from the base of the papillary muscles (*asterisk*) and RV lateral wall (*star*) and tricuspid valve annulus (*diamond*).

Download English Version:

<https://daneshyari.com/en/article/8923866>

Download Persian Version:

<https://daneshyari.com/article/8923866>

[Daneshyari.com](https://daneshyari.com)