



ORIGINAL ARTICLE

Laparoscopic training by use of a physical simulator and its application in the general surgery residency



A.J. Guerrero-Hernández, J.T. Palacios-Zertuche,
F.J. Reyna-Sepúlveda, G.E. Muñoz-Maldonado*

General Surgery Service at the "Dr. José Eleuterio González" University Hospital of the Autonomous University of Nuevo León, Monterrey, Mexico

Received 14 June 2016; accepted 29 September 2016
Available online 16 June 2017

KEYWORDS

Laparoscopic physical simulator;
Laparoscopic training;
Decreased surgical time

Abstract

Introduction: Minimally invasive techniques have become the gold standard for a number of surgical procedures. The benefits include a decrease in morbidity, reducing the length of hospital stay and a faster return to daily activities.

Material and methods: An experimental, longitudinal, comparative, prospective, non-blinded study. A training program of 4 weeks' duration was carried out with an experimental group of first and second year General Surgery residents and social service medical interns (group A); at the end of the training there was a test, where a series of exercises compared the residents who took the training program with a control group (group B) composed of residents from the fourth and fifth year who did not take this training.

Results: Upon analyzing the two groups, we observed a decrease in the average time by group A compared to group B. Overall, the 7 exercises show a difference in average time, with group A completing them in 5 min compared to group B's 12 min ($p < 0.05$).

Discussion: This study shows that training in a physical simulator for laparoscopic surgery, for at least two hours a week for a month, is sufficient to improve laparoscopic surgical skills, reducing turnaround times by more than half compared to those who did not undergo this training.

© 2017 Published by Masson Doyma México S.A. on behalf of Universidad Autónoma de Nuevo León. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Servicio de Cirugía General, Hospital Universitario "Dr. José Eleuterio González" de la Universidad Autónoma de Nuevo León, Avenida Francisco I. Madero y Gonzalitos S/N, Colonia Mitras Centro, Monterrey, C.P. 64460 Nuevo León, Mexico.
Tel.: +52 81 83483303.

E-mail address: cevam99@gmail.com (G.E. Muñoz-Maldonado).

Introduction

Minimally invasive techniques have become the gold standard for a number of surgical procedures. Benefits include a drop in morbidity, thus reducing hospital stay, and a faster return to everyday activities.¹

The term "learning curve" was introduced in surgery to make a reference to the number of surgeries that a surgeon must perform in order to reach a level of experience with a low incidence of the range of complications. Depending on the type of surgery, between 15 and 100 procedures are required to reach the learning curve plateau. Even experienced laparoscopic surgeons have to go through the learning curve again when they train for new laparoscopic techniques or instruments.²

For a long time, surgical teaching and training has been based on a teacher-student model, applied mainly in the operating room. In recent years, however, there have been major changes in surgery. Additionally, the available time for residents in their first years has shortened, and new surgical techniques have emerged. These new techniques lead to new and significant demands in surgical teaching and training, resulting in new methods to acquire experience and skills.

The use of a visual feedback base in monitors or other types of screens gives laparoscopy a wide range of teaching and training possibilities based on computing systems.³

Intracorporeal knot-tying and suturing is amongst the most difficult tasks in laparoscopy, involving hand-eye coordination skills and knowledge of the proper steps. Proficiency in laparoscopic sutures is an important requirement for surgeons who wish to conduct advanced laparoscopy.⁴ A non-randomized study showed a significant improvement in intracorporeal suturing by acquiring skills training in a laparoscopic training box.⁵

There are virtual reality simulators or laparoscopic training boxes used for training,⁶ whose objective is the development of the necessary skills and understanding of the spatial-temporal frame of reference and required maneuvering to manipulate instruments and tissue in a bi-dimensional vision, camera manipulation and the ability to control instruments with both hands; thus being able to perform complicated surgical procedures with minimal invasion.⁷⁻⁹

The objective of this study is to prove that applying a training program using a laparoscopic training box during a general surgeon's residency will result in a shortening of time when performing laparoscopic techniques.

Material and methods

An experimental, longitudinal, comparative, prospective non-blind study was conducted. A 4-week training program was carried out at the Department of General Surgery of a teaching Hospital in northeastern México. The program included an experimental group of first and second-year general surgery residents, as well as interns doing their social service (group A). At the end of the training, a test was conducted where they performed a series of exercises comparing the residents who completed the training program with those in the control group (group B), which

included fourth and fifth-year residents who did not complete said training. The program began with the completion of a laparoscopic skills test in a training box, where 7 exercises were performed, bimanual transference using beads toward a post, beads transfer from one container to another, predetermined cut, traction and counter-traction, simulation of intestine inspection using a string, and points of the suture with intracorporeal and extracorporeal knots. This practice was conducted for at least 2 h a week for 4 weeks. The exercises consisted of the following:

- I. *Intracorporeal knot*: Two suture points will be done with a 4-lacing intracorporeal knot, the total time spent on this knot will be measured. Making an intracorporeal knot on a fixed surface is a common practice in laparoscopic surgery.
- II. *Beads 5/5*: 10 beads will be placed (5 yellow and 5 blue) in a container at the center of the image, and there will be two posts, one on the right and another on the left of the container. Using the dominant hand, 5 beads of the same color will be placed on their opposite post, and with the non-dominant hand the rest of the beads which are a different color will be placed on the opposite post. The complete process will be timed from start to finish. This skill is required to perform fine movements and manage depth, necessary skills for staple placement.
- III. *Beads 10/10*: Transfer 10 beads from one container to another using the dominant hand and return them to the first container using the non-dominant hand, the amount of time required to finish the exercise will be measured. This skill is necessary to perform fine movement and to moderate pressure strength similar to that required to extract lines from the abdominal cavity.
- IV. *Predetermined cut*: A predetermined cut will be made (a circle in a piece of cloth) and the amount of time required to complete the circle will be timed. Skill required in the management of laparoscopic scissors with a single hand and presentation with the other hand is necessary for cuts in laparoscopic surgeries.
- V. *String*: Simulation of intestine inspection using a meter-long string with 3 marks, measuring the amount of time it takes to reach the last one. Tests the ability to use both hands, as specified by the simulating intestine inspection.
- VI. *Grape decortication*: An exercise in tension and strain relief will be made by decortication of a grape of medium size and the time required to completely peel it will be measured. This ability is required for the use of both hands, well-controlled use of force and pressure of the fine movements required for the detachment of ligaments and loose tissues and dissection and exposure of structures.
- VII. *Extracorporeal knot*: A suture will be made with 4 extracorporeal knot loops and the time to finish is measured. Making an extracorporeal knot using the low knots is a common practice in laparoscopic surgery.

Upon completion of the four week training program, participants in the experimental group (group A) must have completed 90% of the previously specified hours (7.2 h) and have performed at least 90% of the indicated time periods. A comparative test was performed where residents of Group

Download English Version:

<https://daneshyari.com/en/article/8926042>

Download Persian Version:

<https://daneshyari.com/article/8926042>

[Daneshyari.com](https://daneshyari.com)