



ORIGINAL ARTICLE

Somatotype of patients with type 2 diabetes in a university hospital in Mexico



K. Urrutia-García*, T.J. Martínez-Cervantes, O. Salas-Fraire, N.P. Guevara-Neri

Department of Sports Medicine and Rehabilitation of the "Dr. José Eleuterio González" University Hospital of the Universidad Autónoma de Nuevo León, Mexico

Received 12 September 2014; accepted 21 January 2015

Available online 13 May 2015

KEYWORDS

Somatotype;
Endomorph;
Mesomorph;
Ectomorph;
Diabetes

Abstract

Objective: The aim of the study was to determine the somatotype of Mexican type 2 diabetes patients, using the Heath and Carter somatotype method.

Methods: The study was conducted on 180 subjects, who underwent an anthropometry following the restricted format established by the International Society for the Advancement of Kinanthropometry (ISAK). A database was elaborated and we obtained descriptive measures such as age, weight, height and the 3 somatotype components.

Results: The average age was 58 years (± 11 SD); 58.6 years (± 10.9 SD) for males and 56.8 years (± 11 SD) for females. The average weight was 77.5 kg (± 16.7); 80.7 kg (± 14.6 SD) for males and 75 kg (± 17.8 SD) for females. The results indicate that the mean somatotype for Mexican type 2 diabetes patients was 6.3, 6.4, 0.6. Diabetic females have higher mean values for endomorphy (7.3) and mesomorphy (6.7) and lower mean values for ectomorphy (0.4) than their male counterparts (5.0, 6.0 and 0.8, respectively). It is evident that endomorphy is predominantly in females, in contrast to males, since there are known differences in fat and muscular mass between both genders.

Conclusions: The results are similar to previous studies presented for other diabetic populations. Sex differences are significant and especially higher for the endomorphic component, with generally higher values in females. It is important to continue this research direction with bigger samples and the study of other risk factors that directly affect the somatotype of type 2 diabetes.

© 2014 Universidad Autónoma de Nuevo León. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Departamento de Medicina del Deporte y Rehabilitación del Hospital Universitario "José Eleuterio González" de la Universidad Autónoma de Nuevo León, Ave. Gonzalitos Col. Mitras Centro, C.P. 64460 Monterrey, Nuevo León, Mexico. Tel.: +52 81 83294207.

E-mail address: k.urrutia@hotmail.com (K. Urrutia-García).

Introduction

Diabetes mellitus is a global health problem, as well as the most well-known disorder for centuries. However, the knowledge of its etiology, natural history and epidemiology remain incomplete.¹⁻² The worldwide prevalence of diabetes mellitus has increased exponentially. In the last three decades, the number of people with type 2 diabetes has doubled.³ Despite being an elderly related disease, in the last few years the number of young people diagnosed with the disease is on the rise. Within its pathophysiology, the genetic component is of great importance in diabetes mellitus; nevertheless, it has been linked to obesity and scarce physical activity or a sedentary lifestyle.⁴

Mexico has suffered the fastest changes ever recorded in the history of food and diet patterns and physical activities resulting in obesity. Diabetes plays a major role, being responsible for almost 14% of all deaths. The increasing prevalence of obesity in children and adults, in addition to the metabolic syndrome, suggests that the situation could have a greater impact in the next few years.⁵ The National Health and Nutrition Survey of 2012 estimated that 9.2% of Mexican adults had already been diagnosed with diabetes mellitus. This reflects an important increase in comparison to the observed results in 2000, where the rate was 5.8%, and 2006, with 7%.⁶

The somatotype is a method used to evaluate, study and comprehend body shape and composition in terms of bone, muscle and fatty tissue dimensions.⁷ First described by Sheldon et al. in 1940, later modified by Heath and Carter in 1967, it was defined as the expression of the current typology of the individual.^{8,9} The first component, endomorphy, refers to the relative fat and thinness. The second component, mesomorphy, refers to the relative musculoskeletal development by size unit. And the third component, ectomorphy, refers to the relative linear aspect, based mostly on the size/weight ratio. Ectomorphy assesses the form and degree of longitudinal distribution of the first and second component.^{7,8}

An important and visually useful tool, individually and for averages, is the somatochart. This should be used on a more routine basis. It is the graphic representation of a two-dimensional format which uses X and Y coordinates from the obtained measurements. The way a somatochart is distributed for a sample provides useful information and should be accompanied by a statistical analysis. The relationship between the somatotype and different diseases was first described by Sheldon et al. in 1940. In 2002, Koleva et al. evaluated the relationship between the somatotype and its main components with the prevalence of several chronic diseases. In five groups of patients, the prevalence was significantly linked to the somatotype.¹⁰

Type 2 diabetes mellitus is a metabolic disorder, and as such it affects the patients' body composition elements. The changes that the disease induces in patients complement its anthropological characteristics. In fact, it is well-known that obesity has a strong positive connection with the development of diabetes mellitus.^{2,11} Tafeit et al. utilized optical equipment (lipometer) to perform subcutaneous fatty tissue measurements in different parts of the body on patients with type 2 diabetes as well as in a control group.

They documented a higher fat percentage in the upper limbs than in the lower limbs, in both male and female.¹²

Several studies have made determinations of the somatotype in patients with type 2 diabetes. In 2013, Baltadijev determined the average somatotype in diabetic Bulgarian women between 40 and 60 years old; observing a mesomorph-endomorph somatotype, differing from the mesomorphic somatotype of the control group. Endomorphy and mesomorphy significantly predominated, while ectomorphy was significantly lower.^{13,14} He also observed and determined the somatotype of the male counterpart within the same age interval and those who presented an endomorph-mesomorph somatotype. Mesomorphy was greater, followed by endomorphy. Ectomorphy was also substantially lower.¹¹

Currently, there are no papers where the somatotype of patients with type 2 diabetes has been established, specifically in the Mexican population. Therefore, the objective of this study was to establish the somatotype of patients with type 2 diabetes of the "Dr. José Eleuterio González" University Hospital in Monterrey, Nuevo León, Mexico, in order to have a phenotype for this type of population and begin a new line of research.

Methods

This study was conducted at the "Dr. José Eleuterio González" University Hospital in Monterrey, Nuevo León, Mexico. The subjects were Mexican patients with a previous diagnosis of type 2 diabetes mellitus selected from the Out Patients Clinics of Internal Medicine, General Medicine, Endocrinology, and Sport Medicine and Rehabilitation wards during the months of December 2013 through April 2014. The patients needed to be willing to have the necessary measurements taken to establish their somatotype. The exclusion criteria were conditions or complications which could alter body composition such as: lower limb edema, amputations or pregnancy. Additionally, other exclusion criteria were situations or conditions which limited their ability to stand up, such as fractures or recent surgeries.

The study included 180 subjects with type 2 diabetes, previously diagnosed. All subjects underwent an anthropometry, under the restrained profile established by the International Society for the Advancement of Kinanthropometry (ISAK).

The ISAK restrained anthropometry profile consists of the taking of base measurements (weight and size), skin folds (triceps, biceps, subscapular, iliac crest, supraspinal, abdominal, anterior thigh, and medial calf), circumferences (relaxed arm, flexed arm, waist, hips and calves) and diameters (bicipital humerus and femur breadth). Regarding skinfolds, circumferences and diameters, these were obtained from the subject's right hemibody as established by ISAK as a standardization method. All measurements were taken by the same measurer in order to avoid variability. Furthermore, in order to reduce variability, the measurer was Level I ISAK certified.

Different equipment was necessary to obtain anthropometric measurements: Fat analyzer TANITA (model TBF-305), to determine weight, and a SECA wall-mounted measuring tape (model 206) to determine size. A body

Download English Version:

<https://daneshyari.com/en/article/8926129>

Download Persian Version:

<https://daneshyari.com/article/8926129>

[Daneshyari.com](https://daneshyari.com)