



Arthroscopic repair of an isolated teres minor tear with associated posterior glenohumeral ligament avulsion

Aaron Schwartz, MD ^{*}, Spero Karas, MD

Sports Medicine, Emory University, Atlanta, GA, USA



ARTICLE INFO

Keywords:

Teres minor tear
Arthroscopic repair
Posterior glenohumeral ligament avulsion
Reverse HAGL
Shoulder arthroscopy
Rotator cuff tear

Isolated teres minor tears are very rare, and only a few case reports have been published in the literature.^{6,7,11,12} The teres minor functions as an external rotator of the glenohumeral joint and a secondary depressor of the humeral head. Most teres minor tears are associated with larger rotator cuff tears; however, there is minimal literature on how to manage isolated injuries. Posterior humeral avulsion of the glenohumeral ligaments (HAGL) is rare as well and is usually associated with labral tears.^{1,3,4,10} Only 1 case report has been published in the literature describing a teres minor avulsion and concomitant bony posterior HAGL injury, although it was also associated with a posterior Bankart lesion.¹² In this case report we describe a new arthroscopic technique for the repair of a teres minor tear with an associated soft tissue posterior HAGL lesion, without other concomitant injuries.

Case report

A 15-year-old right-hand-dominant boy presented with left shoulder pain 10 days after he sustained a traumatic injury while wrestling. He felt his shoulder “shift” with associated pain. He had been using a sling since the injury occurred. On examination, he had full range of motion but had discomfort with active flexion and abduction. He displayed weakness in external rotation but had good strength in the rest of his rotator cuff musculature. He had no specific point tenderness and had no shoulder instability. A magnetic resonance image (MRI) from another facility showed an avulsion of the teres minor from its insertion on the humeral head and a posterior HAGL lesion (Figs. 1 and 2). The patient was sent to physical therapy while a surgical date was set 3 weeks later.




Figure 1 Coronal magnetic resonance imaging study with the arrow pointing to an isolated teres minor tear.

Surgical technique

Arthroscopy was performed with the patient in lateral decubitus. A standard posterior viewing portal was established, followed by an anterior portal in the rotator interval. All intra-articular structures were visualized and inspected with a 30° arthroscope. The biceps, labrum, chondral surfaces, supraspinatus, and infraspinatus were free of injury. The posterior glenohumeral ligament injury was visualized after the arthroscope was put in the anterior portal

The patient and his father provided written permission for this case report.

^{*} Corresponding author: Aaron Schwartz, MD, Emory University, 675 Ponce De Leon Ave NE, Apt W731, Atlanta, GA 30308, USA.

E-mail address: schwaaaron@gmail.com (A. Schwartz).




Figure 2 Axial magnetic resonance imaging study with the arrow pointing to the posterior humeral avulsion of the glenohumeral ligaments avulsion.

(Fig. 3). The arthroscope was then placed into the subacromial space and an accessory anterolateral, accessory posterolateral portal and the port of Wilmington were established (Fig. 4).

A clear 8.25-mm cannula (Arthrex, Naples, FL, USA) was placed in the superior posterolateral portal to allow passage of sutures without creating a tissue bridge. The avulsion of the teres minor had filled in with scar tissue and hematoma. That area was débrided with a shaver (Arthrex) and coblation wand (Smith and Nephew, London, UK), and the anatomic plane was visualized (Fig. 5).

The area beneath the avulsed tendon was prepared using a 5.5-mm burr to create a bony bed for healing. Two FiberTape sutures (Arthrex) were passed in a mattress fashion from distal to proximal along the tear using an ExpressSew (DePuy Synthes, Raynham, MA, USA). This effectively captured the posterior glenohumeral ligament along with the teres minor tendon with the suture. Two 4.75-mm SwiveLock anchors (Arthrex) were placed to reinsert the tendon




Figure 4 Arthroscopic portals: A, standard posterior portal; B, port of Wilmington; C, accessory posterolateral portal; D, accessory anterolateral portal, and E, standard anterior portal.

and the posterior HAGL lesion onto the humeral head with knotless fixation (Fig. 6). A small dog-ear was tied down with a No. 2 FiberWire to reinforce the repair. The arthroscope was then inserted back into the anterior interval portal to view the repair of the HAGL lesion (Fig. 7).

Postoperative outcome

The patient was placed in a sling for 6 weeks and started pendulum range of motion exercises 2 days postoperatively. At 3 weeks, the patient began gentle passive forward elevation with an internal rotation limit to the beltline. Active motion was started at 6 weeks, and strengthening was started after 3 months.

The patient was seen 6 months postoperatively. He was having no pain with activities, had nearly complete range of motion in all planes, and had full external rotation strength. He was told to pro-




Figure 3 Arthroscopic intra-articular view from the anterior portal shows the posterior humeral avulsion of the glenohumeral ligaments injury.



Figure 5 Arthroscopic bursal-sided view from the accessory posterolateral portal shows the teres minor injury.

Download English Version:

<https://daneshyari.com/en/article/8926597>

Download Persian Version:

<https://daneshyari.com/article/8926597>

[Daneshyari.com](https://daneshyari.com)