


Accepted Manuscript

IMPACT OF SUBSTANCE USE DISORDER ON GRAY MATTER VOLUME IN SCHIZOPHRENIA

Margaret Quinn , Maureen McHugo , Kristan Armstrong ,
Neil Woodward , Jennifer Blackford , Stephan Heckers

PII: S0925-4927(18)30004-0
DOI: <https://doi.org/10.1016/j.psychresns.2018.08.002>
Reference: PSYN 10845



To appear in: *Psychiatry Research: Neuroimaging*

Received date: 12 January 2018
Revised date: 14 May 2018
Accepted date: 2 August 2018

Please cite this article as: Margaret Quinn , Maureen McHugo , Kristan Armstrong , Neil Woodward , Jennifer Blackford , Stephan Heckers , IMPACT OF SUBSTANCE USE DISORDER ON GRAY MATTER VOLUME IN SCHIZOPHRENIA, *Psychiatry Research: Neuroimaging* (2018), doi: <https://doi.org/10.1016/j.psychresns.2018.08.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Substance use may confound the study of brain structure in schizophrenia.
- We compared schizophrenia patients with and without substance use histories.
- We replicated gray matter volume deficits known to occur in schizophrenia.
- These did not significantly differ in patients with and without substance use.
- A substance use history does not compound gray matter deficits in schizophrenia.

Download English Version:

<https://daneshyari.com/en/article/8941698>

Download Persian Version:

<https://daneshyari.com/article/8941698>

[Daneshyari.com](https://daneshyari.com)