



Contents lists available at ScienceDirect

Schizophrenia Research

journal homepage: [www.elsevier.com/locate/schres](http://www.elsevier.com/locate/schres)

## Acoustic startle response and its modulation in schizophrenia and autism spectrum disorder in Asian subjects

Hidetoshi Takahashi<sup>a,b,\*</sup>, Yoko Kamio<sup>a</sup>

<sup>a</sup> Department of Child and Adolescent Mental Health, National Institute of Mental Health, National Center of Neurology and Psychiatry, 4-1-1 Ogawahigashicho, Kodaira, Tokyo 187-8553, Japan

<sup>b</sup> Department of Advanced Neuroimaging, Integrative Brain Imaging Center, National Center of Neurology and Psychiatry, 4-1-1 Ogawahigashicho, Kodaira, Tokyo 187-8551, Japan

### ARTICLE INFO

#### Article history:

Received 27 February 2017

Received in revised form 23 May 2017

Accepted 26 May 2017

Available online xxxx

#### Keywords:

Acoustic startle response

Prepulse inhibition

Sensorimotor gating

Schizophrenia

Autism spectrum disorder

Asian population

### ABSTRACT

The acoustic startle response (ASR) and its modulation, including prepulse inhibition (PPI), are considered to be promising neurophysiological indices for translational research in psychiatry. Impairment of the PPI has been reported in several psychiatric disorders, but particularly in schizophrenia, where PPI is considered to be a candidate endophenotype of the disorder. Although the profiles of the ASR differ between races, recent studies of single ethnicity samples in Asia were in accord with a number of studies from Western countries, in reporting that patients with schizophrenia exhibit impaired PPI.

The PPI of the ASR is known to develop before 8 years of age, and PPI impairment has only been reported in adults (not children) with autism spectrum disorder (ASD), which involves atypical features that are present from early development. Recent Asian studies of children with ASD suggest that comprehensive investigation of the ASR and its modulation, including the startle response to weak startle stimuli, peak startle latency, and PPI, may contribute to an understanding of the impairment of the neural circuitry in children with ASD and its comorbid behavioral problems.

In this review, we review recent findings on the ASR and its modulation from Asian countries, and discuss its potential use for studying sensorimotor gating and its relationship to schizophrenia and ASD.

In conclusion, the ASR and its modulation can provide a well-established global neurophysiological index for translational research in psychiatric disorders. Future studies investigating the development of sensorimotor gating in early development may contribute to prevention of psychiatric disorders.

© 2017 Elsevier B.V. All rights reserved.

### 1. Introduction

The acoustic startle response (ASR) and its modulation, including prepulse inhibition (PPI), are considered to be some of the most promising neurophysiological indices for translational research in psychiatry; research that combines both basic and clinical research for the purpose of understanding the complex pathogenesis of the genetic and environmental interactions underlying psychiatric disorders. As the ASR can be assessed using simple nonlinguistic stimuli, the ASR, and especially the related PPI, have been widely investigated across human races (Hasenkamp et al., 2008; Swerdlow et al., 2007, 2005) and other species (Powell et al., 2012, 2009; Swerdlow and Geyer, 1998; Swerdlow and Light, 2016) using similar experimental paradigms (Light and Swerdlow, 2014; Swerdlow and Light, 2016).

The ASR is a fast twitch of facial and body muscles in response to a sudden and intense sound (Koch, 1999). This response pattern suggests that the startle may have a role in protecting against injury from a predator or a blow, and is an element of the flight response. PPI is usually defined as a reduction in the startle reflex due to weak sensory prestimulation (Braff et al., 1978), and is the most commonly used psychophysiological index of sensorimotor gating, which is an autonomic inhibition system that regulates the motor response to sensory input. This inhibitory process has been proposed to contribute to the selective and efficient processing of relevant information.

Impairment of the PPI has been reported in several psychiatric disorders (Swerdlow et al., 2016; Takahashi et al., 2011), of which schizophrenia is the most prominent. Although the profile of the ASR differs across racial groups (Hasenkamp et al., 2008; Swerdlow et al., 2007, 2005), studies of both Asian and Caucasian subjects indicate that patients with schizophrenia exhibit impaired PPI, and that impaired sensorimotor gating may be a global common psychophysiological feature of schizophrenia.

Investigation of the ASR and its modulation in other neurodevelopmental disorders, such as autistic spectrum disorder (ASD; McAlonan et al., 2002; Ornitz et al., 1993; Perry et al., 2007;

\* Corresponding author at: Department of Child and Adolescent Mental Health, National Institute of Mental Health, National Center of Neurology and Psychiatry, 4-1-1 Ogawahigashicho, Kodaira, Tokyo 187-8553, Japan.

E-mail addresses: [htakahashi@ncnp.go.jp](mailto:htakahashi@ncnp.go.jp) (H. Takahashi), [kamio@ncnp.go.jp](mailto:kamio@ncnp.go.jp) (Y. Kamio).

Takahashi et al., 2016; Yuhas et al., 2011) or Tourette's syndrome (Swerdlow, 2013; Swerdlow et al., 2001), may provide important insights into the underlying mechanism behind the developmental aspect of sensorimotor gating and its relationship to psychiatric disorders. In particular, ASD, which has atypical features from early development, is known to have a relatively high incidence (12–17%) of comorbid psychotic disorders, which manifest mainly in adults (Lai et al., 2014). Although the developmental timing of symptom onset clearly differs between schizophrenia and ASD, there are several similarities in their clinical presentation, including social dysfunction and sensory abnormalities (Konstantareas and Hewitt, 2001), as well as genetics (State and Levitt, 2011) and neurobiology (State and Sestan, 2012). Thus, basic investigation of the ASR and its modulation in children with ASD may provide deeper insight into the development of sensorimotor gating, which is related to psychotic disorders.

In this review, we review recent findings related to the PPI of the ASR in schizophrenia from a single ethnicity Asian country; findings that suggest that impaired sensorimotor gating is a global common psychophysiological feature of schizophrenia. We also review recent studies on ASR and its modulation in children with ASD; an area of research that is evolving in Asian countries, particularly for the purpose of forwarding the understanding of the developmental aspects of auditory sensory processing in relation to psychiatric disorders.

## 2. Prepulse inhibition of the ASR in Asian patients with schizophrenia

Schizophrenia is one of the most prominent psychiatric disorders presenting with deficits in PPI. Impaired sensorimotor gating has been considered to be a common psychophysiological feature of schizophrenia, which may theoretically lead to severe dysfunctions in perception, attention, and thinking (Braff and Geyer, 1990). Since the first report on their occurrence (Braff et al., 1978), reductions in PPI have been consistently demonstrated in patients with schizophrenia (Braff et al., 2001a, 2001b; Swerdlow et al., 2014, 2008), and recently, PPI has been

considered a promising candidate endophenotype of schizophrenia (Braff et al., 2007; Braff and Light, 2005; Greenwood et al., 2007, 2016, 2013; Swerdlow et al., 2014; Turetsky et al., 2007).

Reduced PPI in patients with schizophrenia has also been consistently reported in recent studies on Asian subjects of single ethnicities (Kunugi et al., 2007; Moriwaki et al., 2009; Takahashi et al., 2008), despite the smaller magnitude of the ASR in Asian subjects in comparison to Caucasians (Swerdlow et al., 2007, 2005). Table 1 shows a summary of previous ASR studies of schizophrenia and related conditions in Asian populations. Most PPI studies on Asian subjects with schizophrenia have reported on single ethnicity Japanese patient samples. Firstly, a Japanese study of PPI in schizophrenia (Kunugi et al., 2007) examined the PPI of the ASR in 20 Japanese patients with chronic schizophrenia under antipsychotic medication, and compared it with that in 16 healthy Japanese controls matched for age and sex. The PPI differed significantly between the two groups. Later Japanese studies with relatively larger sample sizes (Kishi et al., 2012, 2013; Matsuo et al., 2016; Moriwaki et al., 2009; Takahashi et al., 2008) have also consistently reported a reduction of PPI in patients with schizophrenia (n: 51–181) in comparison with healthy controls (n: 55–250). Some of these studies also reported reduced habituation of the ASR in Japanese patients with schizophrenia (Matsuo et al., 2016; Moriwaki et al., 2009). Habituation is another ASR modulation that is usually defined as a decrease in the behavioral response after repeated presentation of an identical stimulus, which is not caused by sensory adaption or effector fatigue (Geyer and Braff, 1982). Reduced ASR habituation may reflect the impaired gating of repeated stimuli, and may result in cognitive disruption via sensory overload. Reduced PPI of the ASR in schizophrenia have also been observed in Chinese samples (Wang et al., 2012; Xue et al., 2012). In Chinese samples, PPI was found to be smaller in both medicated and unmedicated patients with schizophrenia, compared with healthy controls (Shi et al., 2016). Although the majority of studies of Asian patients have reported a reduction of ASR and its habituation in patients with schizophrenia (Table 1), these features are not consistently observed in Caucasian patients with schizophrenia (Braff et al., 2001a, 2001b;

**Table 1**  
Summary of studies on acoustic startle response and its modulation in Asian patients with schizophrenia.

Study	SCZ	Age [years] (mean ± SD)	HP	Age [years] (mean ± SD)	PI (dB SPL)	LI (msec)	ITI (sec)	Main findings
Kunugi et al., 2007	20 (chronic) Male 12	42 ± 9	16 Male 9	41 ± 13	82, 86, 90	30, 60, 120	10–20 Average 15	ASR, PPI at 120 msec LI smaller in SCZ than in HP.
Takahashi et al., 2008	51 Male 29	38.1 ± 10.2	55 Male 28	40.4 ± 10.1	82, 86, 90	120	15–25 Average 20	HAB, PPI for all prepulse intensities smaller in SCZ than in HP.
Moriwaki et al., 2009	111 (chronic) Male 71	52.0 ± 1.46	110 Male 72	32.7 ± 0.674	82, 86, 90	120	15–25 Average 20	ASR, HAB, PPI for all prepulse intensities smaller in SCZ than in HP.
Wang et al., 2012	50 (chronic) Male only	data not shown	25 Male only	data not shown	85	30, 60, 120	11–19 Average 15	ASR, HAB, PPI at 120 msec LI smaller in SCZ taking typical antipsychotics compared to HP (25 treated with typical antipsychotics, 25 with clozapine).
Wang et al., 2013	31 Male 17	20.4 ± 4.5	30 Male 13	22.1 ± 3.5	85	30, 60, 120	11–19 Average 15	ASR, PPI at 60 msec LI smaller in SCZ than in HP (first-episode, medication-naïve).
Kishi et al., 2013	100 (chronic) Male 66	50.8 ± 15.5	107 Male 71	32.7 ± 7.16	82, 86, 90	120	15–25 Average 20	ASR, HAB, PPI for all PIs smaller in SCZ than in HP.
Matsuo et al., 2016	181 Male 83	37.6 ± 11.2	250 Male 125	36.7 ± 12.4	86, 90	60, 120	10–20 Average 15	ASR, HAB, PPI at 90 dB PI and 60, 120 msec LI smaller in SCZ than in HP.
Kishi et al., 2010	141 (chronic) Male 91	49.8 ± 15.6			82, 86, 90	120	15–25 Average 20	PPI at 90 dB SPL PI was larger in SCZ taking aripiprazole or risperidone compared to those taking olanzapine.
Kishi et al., 2012	100 (chronic) Male 66	50.8 ± 15.5			82, 86, 90	120	15–25 Average 20	HAB was associated with level of attention in SCZ.

Background noise intensity 70 dB SPL, pulse duration 40 msec, and pulse intensity 115 dB SPL, prepulse duration 20 msec in all studies. ASR: acoustic startle response; HAB: habituation; HP: healthy participants; ITI: Inter-trial intervals; LI: lead interval from prepulse onset to pulse onset; PI: prepulse intensity; PPI: prepulse inhibition of ASR; SCZ: patients with schizophrenia; SPL: sound pressure level.

Download English Version:

<https://daneshyari.com/en/article/8941710>

Download Persian Version:

<https://daneshyari.com/article/8941710>

[Daneshyari.com](https://daneshyari.com)