

Accepted Manuscript

Preoperative smoking, narcotic, benzodiazepine and tramadol use are risk factors for narcotic use after hip and knee replacement

Kipp A. Cryar, MD, Timothy Hereford, Paul K. Edwards, MD, Eric Siegel, C.Lowry Barnes, MD, Simon C. Mears, MD



PII: S0883-5403(18)30345-0

DOI: [10.1016/j.arth.2018.03.066](https://doi.org/10.1016/j.arth.2018.03.066)

Reference: YARTH 56560

To appear in: *The Journal of Arthroplasty*

Received Date: 12 February 2018

Revised Date: 21 March 2018

Accepted Date: 30 March 2018

Please cite this article as: Cryar KA, Hereford T, Edwards PK, Siegel E, Barnes CL, Mears SC, Preoperative smoking, narcotic, benzodiazepine and tramadol use are risk factors for narcotic use after hip and knee replacement, *The Journal of Arthroplasty* (2018), doi: 10.1016/j.arth.2018.03.066.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Preoperative smoking, narcotic, benzodiazepine and tramadol use are risk factors for narcotic use
after hip and knee replacement

Kipp A. Cryar, MD

Timothy Hereford

Paul K. Edwards, MD

Eric Siegel

C. Lowry Barnes, MD

Simon C. Mears, MD

Department of Orthopaedic Surgery, University of Arkansas for Medical Sciences, Little Rock,
AR, USA.

Please address all correspondence to:

Simon C. Mears, MD PhD

4301 West Markham Street, Slot 531

Little Rock, AR 72205

Tel/Fax: 501-686-6067

E-mail: SCMears@uams.edu

Download English Version:

<https://daneshyari.com/en/article/8945447>

Download Persian Version:

<https://daneshyari.com/article/8945447>

[Daneshyari.com](https://daneshyari.com)