Accepted Manuscript



Multiple-Dose Intravenous Tranexamic Acid Further Reduces Hidden Blood Loss After Total Hip Arthroplasty: A Randomized Controlled Trial

Yiting Lei, MM, Qiang Huang, MD, Zeyu Huang, MD, PhD, Jinwei Xie, MD, Guo Chen, MD, Fuxing Pei, MD

PII: S0883-5403(18)30385-1

DOI: 10.1016/j.arth.2018.04.024

Reference: YARTH 56588

To appear in: *The Journal of Arthroplasty*

Received Date: 24 January 2018

Revised Date: 13 March 2018

Accepted Date: 15 April 2018

Please cite this article as: Lei Y, Huang Q, Huang Z, Xie J, Chen G, Pei F, Multiple-Dose Intravenous Tranexamic Acid Further Reduces Hidden Blood Loss After Total Hip Arthroplasty: A Randomized Controlled Trial, *The Journal of Arthroplasty* (2018), doi: 10.1016/j.arth.2018.04.024.

Multiple-Dose Intravenous Tranexamic Acid Further Reduces Hidden Blood Loss After Total Hip Arthroplasty: A Randomized Controlled Trial

Yiting Lei, MM [a]; Qiang Huang, MD [a]; Zeyu Huang, MD, PhD [a]; Jinwei Xie, MD [a]; Guo Chen, MD [a]; Fuxing Pei, MD [a]

Corresponding Author: Fuxing Pei, E-mail: peifux@126.com; Telephone: 13258327318

[a] Department of Orthopaedics, West China Hospital, Sichuan University, 37# WainanGuoxue Road, Chengdu 610041, People's Republic of China

Yiting Lei and Qiang Huang contributed equally to this work and should be considered as equal first authors.