

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Transfer of the anterior branch of the obturator nerve for femoral nerve reconstruction and preservation of motor function: A case report

Marco Rastrelli^{a,1}, Ilaria Tocco-Tussardi^{b,*,1}, Saveria Tropea^a, Carlo Riccardo Rossi^a, Sandro Rizzato^b, Vincenzo Vindigni^b

^a Istituto Oncologico Veneto [Veneto Institute of Oncology], Via Gattamelata 64, 35128 Padova, Italy

^b Clinic of Plastic and Reconstructive Surgery, Department of Neurosciences, University of Padova, Via Giustiniani 2, 35128 Padova, Italy

ARTICLE INFO

Article history:

Received 17 May 2018

Received in revised form 19 July 2018

Accepted 2 August 2018

Available online 9 August 2018

Keywords:

Anterior branch of the obturator nerve

Femoral nerve

Gracilis

Nerve transfer

Neurotization

ABSTRACT

INTRODUCTION: Femoral nerve lesions are uncommon but severely disrupting at the functional level, because of the inability to walk, run, and passing from sitting to standing position. Reconstruction via local nerve transfer (neurotization) is a relatively new yet promising procedure.

PRESENTATION OF CASE: We describe a case of successful restoration of rectus femoris' function after a malignant process by transfer of the anterior branch of the obturator nerve at the thigh level. At present, only few similar reports are present in the literature. Nerve gap after surgery was considerable (10 cm) and nerve grafting could have been unsatisfactory in terms of reinnervation. Therefore, reconstruction was managed with nerve isolation and transfer to the rectus femoris motor branch. The functional result was satisfactory at 1-year follow up with margins for further improvement.

DISCUSSION: This case reported favorable outcomes of neurotization of the anterior branch of the obturator nerve for femoral nerve lesion. Reports of success with this procedure are still limited, but the promising results in terms of functional recovery suggest it should be offered to patients as a viable therapeutic option.

CONCLUSION: Advantages of neurotization compared to grafts are several, including: limiting suturing sites and scarring; shortening the recovery time by decreasing the required regeneration distance; and allowing for faster muscle reinnervation. The choice to transfer the anterior branch of the obturator nerve specifically allows to preserve part of the adductor functionality in the thigh without affecting the stability of the knee joint.

© 2018 The Author(s). Published by Elsevier Ltd. on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Damage to the femoral nerve and subsequent nerve palsy can occur as a consequence of trauma (pelvic fractures, gun-shot wounds), surgical procedures (intra-abdominal, vascular, gynecological surgery), or as an outcome of nerve sheath tumor (Schwannoma, neurofibroma, neurosarcoma)/infiltrating tumor surgery [1,2]. The resulting paralysis of the femoral quadriceps causes significant disability, with a severe impairment and reduction of the quality of life. In cases of a significant nerve gap, nerve transfer (neurotization) can be considered as a valid therapeutic option, to restore nerve continuity in order to re-donate function. We describe a case of successful restoration of rectus femoris' function after a malignant process by transfer of the anterior branch of

the obturator nerve at the level of the thigh. The case has been reported in line with the SCARE criteria [3].

2. Presentation of case

A 19-year-old girl was referred to the Multidisciplinary Sarcoma Group for evaluation from another Hospital, where three months before she had undergone excision of a 'myxoid' liposarcoma located on the anterior aspect of the right upper thigh. From pre-operative RM study, the tumor was described as a 'solid, expansive, lobulated, septated mass starting from the groin, bordered by sartorius and iliopsoas muscles, causing no alterations in the common femoral artery (which for 50% of its diameter was bordered by the lesion) and vein normal permeability', with no changes in bone cortex of the right femur, and no inguinal lymphadenopathy; dimensions were 2.6 × 6.4 × 9.5 cm. Excision had been performed with an incision under the inguinal ligament, and a drainage tube had been kept in place for four weeks. His-

* Corresponding author.

E-mail address: ilaria.tocottussardi@gmail.com (I. Tocco-Tussardi).

¹ These authors contributed equally to this work.



Fig. 1. Intra-operative photographs. (Left) Operative exposure. (Right) After transposition of the anterior branch of the obturator nerve to femoral branches to rectus femoris, the soft tissue defect is covered with a rotation of muscular flap.

tological report had revealed the liposarcoma had been excised by fragmentation. Molecular cytogenetic analysis by fluorescence in situ hybridization (FISH) had revealed gene rearrangements involving CHOP/DDIT3, supporting diagnosis. The patient had not received adjuvant radiotherapy or chemotherapy. She was otherwise healthy, had multiple allergies (cefotaxime, ceftriaxone, cefazolin, and beta-lactams), and was receiving no drug therapy; she had a positive family history of lung cancer.

A new contrast-enhanced RM scan was obtained, showing a nodular formation on tensor fascia lata muscle. The Multidisciplinary team agreed on a treatment plan of surgical revision of the groin area, with en-bloc removal of the muscles of Scarpa's triangle, the superficial and deep femoral artery and vein, and the femoral nerve.

Options for reconstruction included: exploration and nerve grafting; and nerve transfer more distally in the thigh. We estimated a final nerve gap of 9 cm, starting 1 cm under the inguinal ligament, after the 'en-bloc' removal (skin and subcutaneous tissue, fascia, sartorius, pettineus, and vastus lateralis muscle, femoral artery and vein – common, superficial and deep, and femoral nerve). Therefore, we were concerned that nerve grafting would have been unsatisfactory in terms of reinnervation. Among the options in proximity to the femoral nerve that would allow a direct transfer, the obturator nerve innervating the adductor muscle group was the most feasible potential donor nerve. Nerve transfer was performed from the anterior branch of the obturator nerve, which was isolated in a distal-to-proximal fashion up to the obturator foramen (10 cm), leaving intact the posterior branch to preserve some adductor function. The nerve was transferred to the rectus femoris motor branch of the femoral nerve, under the adductor longus muscle. Femoral arteries were reconstructed with a vein graft (great saphenous vein harvested from the contralateral lower limb), and soft tissue defect was covered with a rotational muscular flap (tensor fasciae latae muscle) and split-thickness skin grafts harvested from the anterolateral aspect of the contralateral thigh (Fig. 1). The post-operative period was uneventful. The histological examination was negative.

At one year post-operatively, there was clear evidence of active leg extension and hip flexion (Fig. 2). She had difficulties running and walking up stairs, but was continuing to improve over time.



Fig. 2. Post-operative photographs 1 year after surgery.

3. Discussion

Neurotization is a well-known reconstructive technique for proximal nerve injuries, with applications documented especially for brachial plexus lesions reconstruction [4–7]. With regard to lower limb, neurotization of the anterior branch of the obturator nerve – motor nerve to the gracilis muscle (MNG) for femoral nerve reconstruction is a recently-acquired technique that is now starting

Download English Version:

<https://daneshyari.com/en/article/8945975>

Download Persian Version:

<https://daneshyari.com/article/8945975>

[Daneshyari.com](https://daneshyari.com)