The insertion of the non-heme FeB cofactor into nitric oxide reductase from P. denitrificans depends on NorQ and NorD accessory proteins

Maximilian Kahle, Josy ter Beek, Jonathan P. Hosler, Pia Ädelroth

PII: S0005-2728(18)30138-5
DOI: doi:10.1016/j.bbabio.2018.05.020
Reference: BBABIO 47932

To appear in: *BBA - Bioenergetics*

Received date: 16 February 2018
Revised date: 27 May 2018
Accepted date: 31 May 2018

Please cite this article as: Maximilian Kahle, Josy ter Beek, Jonathan P. Hosler, Pia Ädelroth , The insertion of the non-heme FeB cofactor into nitric oxide reductase from P. denitrificans depends on NorQ and NorD accessory proteins. Bbabio (2018), doi:10.1016/j.bbabio.2018.05.020

The insertion of the non-heme Fe_B cofactor into Nitric Oxide Reductase from *P. denitrificans* depends on NorQ and NorD accessory proteins

Maximilian Kahle[1], Josy ter Beek[1#], Jonathan P. Hosler[2] and Pia Ädelroth[1*]

[1]Department of Biochemistry and Biophysics, Stockholm University, SE-106 91 Stockholm, Sweden; [2]Department of Biochemistry, University of Mississippi Medical Center, Jackson, Mississippi 39216, United States

[#]Present address: Department of Medical Biochemistry and Biophysics, Umeå University, SE-901 87 Umeå, Sweden

[*]To whom correspondence should be addressed. Pia Ädelroth: Department of Biochemistry and Biophysics, Stockholm University, SE-106 91 Stockholm, Sweden; pia.adelroth@dbb.su.se; Tel. +46-8-164183.

Keywords: iron, AAA ATPases, *c*NOR, metal ion insertion, *nor* accessory genes, His-tag, *E. coli*, chaperone, protein assembly, MoxR