Cannabidiol restores differentiation capacity of LPS exposed adipose tissue mesenchymal stromal cells

Tim Ruhl, Bong-Sung Kim, Justus P. Beier

www.elsevier.com/locate/yexcr

PII: S0014-4827(18)30431-2
DOI: https://doi.org/10.1016/j.yexcr.2018.07.030
Reference: YEXCR11134

To appear in: *Experimental Cell Research*

Received date: 9 July 2018
Revised date: 17 July 2018
Accepted date: 19 July 2018

Cite this article as: Tim Ruhl, Bong-Sung Kim and Justus P. Beier, Cannabidiol restores differentiation capacity of LPS exposed adipose tissue mesenchymal stromal cells, *Experimental Cell Research,* https://doi.org/10.1016/j.yexcr.2018.07.030

Cannabidiol restores differentiation capacity of LPS exposed adipose tissue mesenchymal stromal cells

Tim Ruhl*[1], Bong-Sung Kim[2], Justus P. Beier[3]

Department of Plastic Surgery, Hand Surgery–Burn Center, University Hospital RWTH Aachen, Pauwelsstraße 30, 52074 Aachen, Germany

truhl@ukaacken.de

bkim@ukaachen.de

jbeier@ukaachen.de

***Corresponding author**: Tim Ruhl, PhD, M.Sc. Department of Plastic Surgery, Hand Surgery–Burn Center, University Hospital RWTH Aachen, Pauwelsstraße 30, 52074 Aachen, Germany. Phone: +49 241 80 37428; Fax +49 241 80 37428. e-mail: truhl@ukaachen.de

Abstract

Multipotent mesenchymal stromal cells (MSCs) support wound healing processes. These cells express toll-like receptors (TLRs). TLRs perform important key functions when the immune system is confronted with danger signals. TLR ligation by lipopolysaccharides (LPS) activates MSCs and induces intracellular signaling

[1] 49-241-8037428, 49-241-8037428
[2] 49-241-8089759, 49-241-8037428
[3] 49-241-8089700, 49-241-8037428