




Contents lists available at ScienceDirect

Clinical Nutrition ESPEN

journal homepage: <http://www.clinicalnutritionespen.com>

Original article

Fruit and vegetables in hypertensive women with asymptomatic peripheral arterial disease

Anna Vittoria Mattioli ^{a, *}, Coppi Francesca ^b, Migaldi Mario ^a, Farinetti Alberto ^a^a Department of Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Italy^b Cardiology Division, Azienda Ospedaliera Universitaria, Modena, Italy

ARTICLE INFO

Article history:

Received 20 May 2018

Accepted 26 May 2018

Keywords:

Fruit
Vegetables
Antioxidants
Women

SUMMARY

Background and aims: Fruit and vegetables are considered a very healthy diet useful in the prevention of cardiovascular disease. The present study aims to evaluate intake of fruit and vegetables in hypertensive women and its correlation with asymptomatic atherosclerosis.

Methods and results: A group of 237 women with hypertension was evaluated. Fruit and vegetables consumption were assessed by a self-administered food frequency validated questionnaire completed by an interviewer administered 24 h diet recall. They all underwent ABI.

ABI measurement observed that fruit consumption was inversely associated with pre-clinical atherosclerosis suggesting a protective effect, moreover this association was stronger for vegetables. Increasing intake of vegetables was associated with a lower risk of asymptomatic PAD.

Conclusions: Women with a high intake of fruit and vegetables showed less instrumental sign of pre-clinical peripheral atherosclerosis. Can be suggests that fruit and vegetables play an important role in prevention of atherosclerosis in pre-menopausal women.

© 2018 European Society for Clinical Nutrition and Metabolism. Published by Elsevier Ltd. All rights reserved.

Background

Dietary factors play an important role in the development of CVD and its risk factors and may contribute to the geographic variability in CVD morbidity and mortality [1].

A low consumption of fruit and vegetables (less than 400 g [g] per day) is thought to be one of the top 10 risk factors for mortality and is estimated to result in 1.7 million global deaths a year [1]. Several population-based studies have shown the beneficial effects of high fruit and vegetable consumption on cardiovascular prognosis outcomes [2,3]. The PREDIMED study showed a reduction in PAD associated with the Mediterranean diet [2]. Similarly we reported that women with high adherence to Med D are less likely to developed preclinical atherosclerosis compared to women with a low adherence [3] The ankle-

brachial index (ABI) is a symptom-independent tool that can be used reliably to evaluate asymptomatic pre-clinical atherosclerosis [4].

The present study aims to evaluate intake of fruit and vegetables in hypertensive women and its correlation with asymptomatic atherosclerosis.

Methods

A retrospective analysis on a group of 650 women (age range 45–54 years) was performed. Patients were referred to our clinic from general practitioners for screening and prevention of CVD. We selected women only if they were free of symptoms of PAD, had ABI evaluation and a complete nutritional assessment. We excluded participants with a previous history of cardiovascular disease, (ischemic heart disease, heart failure and stroke), those who did not complete questionnaire, and those who did not undergo ABI evaluation and who did not sign the consent. (Figure 1 supplemental material). From the initial group we analyzed data from 237 women with hypertension [5].

Study was approved by the Local Ethical Review Board and participants signed an informed consent.

* Corresponding author. Department of Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Via del pozzo, 71 41100 Modena, Italy. Fax: +39 59 4224323.

E-mail address: annavittoria.mattioli@unimore.it (A.V. Mattioli).

Data collection

Nutritional status was assessed by measuring weight, body mass index (BMI), waist circumference and waist-to-hip ratio [3].

The prevalence of nutritional parameters was assessed by a self-administered food frequency questionnaire (FFQ) with 116 items and completed by an interviewer-administered 7-day diet recall questionnaire on the day of first visit [3].

The food list in the FFQ was Italianized, and foods commonly eaten in the Emilia Romagna region of Italy were added. For each food class color photographs of three different portions were displayed. Portion sizes were chosen according to many years of experience in dietary surveys in various parts of Italy [3].

Food frequency was evaluated using three categories: daily, weekly and monthly and from 1 to 6 number of times (i.e.: once a day, 3 times a week) and was integrated with specific questions on changes in nutrition habits and lifestyle within the last year. The Mediterranean score was calculated according to Panagiotakos [3]. We assessed fruit and vegetables intake. V/F were separated into colors based on their nutritional density: (1) green (vegetables rich in folic acid such as broccoli); (2) orange (V/F rich in beta-carotene such as carrots); (3) purple (vegetables rich in potassium and folic acid such as red cabbage); (4) yellow (fruits rich in vitamin C such as oranges); (5) red (fruits rich in potassium and vitamin C such as cranberries). Because of inconsistent classification of legumes as a subset of vegetables, we analyze total fruits and vegetables without legumes [6].

A dietician, using a database system, computed food and nutrient intakes from FFQ and dietary recall. The nutrient database was compiled from food composition tables. Smoking and physical activity were also investigated.

ABI measurement

All patients underwent ABI measurement. The ABI is a simple, noninvasive test, measuring the SBP from both brachial arteries and from both the dorsalis pedis and posterior tibial arteries after the

patient has been at rest in the supine position for 10 min by using a Doppler device. The ABI of each leg is calculated by dividing the higher of the dorsalis pedis pressure or posterior tibial pressure by the higher of the right or left arm blood pressure [4].

Statistical analysis

SPSS, V.21.0.1 (SPSS Inc, Chicago, Ill) was used for statistical analysis. Results are presented as mean \pm SD or frequency expressed as a percentage.

To study the association of consumption of fruits and vegetables with PAD, we used Cox proportional hazards regression analysis. For these analyses, consumption of fruits and vegetables were used as continuous and categorical (0–1; 2; ≥ 3 servings per day) variables, in order to obtain the best fitting model. We performed analyses in which we first adjusted for age, BMI, smoking, physical activity, and adherence to Mediterranean Diet (model 1); and additionally for total cholesterol, HDL cholesterol, systolic blood pressure, and hs-CRP (model 2).

In interaction analysis, P interaction < 0.05 was considered to indicate effect modifiers on the association of fruit and vegetable consumption with ABI.

Results

The mean MedD Score was 32.6 ± 3.3 (median score was 30.9). Antioxidants intake was higher in patients with a greater adherence to Med D and was mainly related to fruit and vegetables and in a lower percentage from wine and coffee. Mean consumption of fruits and vegetables were 1.8 ± 1.2 servings/day and 2.5 ± 0.8 serving/day, respectively. Mean plasma vitamin C concentration was 42.6 ± 15.2 $\mu\text{mol/L}$.

Table 1 shows association between clinical characteristics and fruit and vegetable consumption (linear regression analysis).

Table 2 illustrates the association of fruit and vegetables consumption with ABI value after adjustments for age, BMI, smoking, physical activity, and adherence to Mediterranean Diet (model 1)

Table 1
Clinical characteristics of patients and association with fruit and vegetable consumption (linear regression analysis).

Clinical characteristics		Fruit consumption B coeff	P value	Vegetables consumption B coeff	P value
Mean age (years)	54.6 \pm 4.2	0.07	ns	0.08	ns
Weight (Kg)	82.3 \pm 4.3	0.14	0.05	0.097	Trend to 0.05
Body mass index (mean)	28.1 \pm 4.8	0.11	0.05	0.09	ns
Waist circumference (cm)	89.5 \pm 3.3	0.03	ns	0.05	ns
Systolic blood pressure, mmHg	154.5 \pm 7.3	0.093	ns	0.097	Trend to 0.05
Diastolic blood pressure, mmHg	92.4 \pm 8.1	0.065	ns	0.067	ns
Smoking (cigarettes/day)	12 \pm 8	0.11	0.05	0.12	0.01
Alcohol (abstainer) number of pts	25	0.02	ns	0.04	ns
hs-CRP, mg/L (available in 125 pts)	2.3 \pm 1.4	0.05	ns	0.07	ns

Table 2
Association of Fruit and Vegetable consumption with asymptomatic PAD in adjusted model.

PAD	Categories of Fruit 0–1 serving/day references	Categories of Fruit 2 serving/day HR (95% CI)	Categories of Fruit >3–4 cups/day HR (95% CI)	Fruit consumption serving days HR (95% CI)
Model 1	1	0.87 (0.37–1.13)	0.82 (0.45–1.23)	0.82 (0.44–0.8)*
Model 2	1	0.96 (0.44–1.17)	0.95 (0.32–1.9)	0.95 (0.45–1.2)
	Categories of Veget. 0–1 serving/day references	Categories of Veget. 2 serving/day HR (95% CI)	Categories of Veget. >3–4 cups/day HR (95% CI)	Veget consumption serving days HR (95% CI)
Model 1	1	0.82 (0.21–1.1)	0.74 (0.32–0.9)	0.81 (0.44–0.85)**
Model 2	1	0.88 (0.49–1.2)	0.77 (0.45–0.83)	0.83 (0.36–0.92)**

Legend: *p < 0.05; **p < 0.01.

Download English Version:

<https://daneshyari.com/en/article/8950669>

Download Persian Version:

<https://daneshyari.com/article/8950669>

[Daneshyari.com](https://daneshyari.com)