

REVISTA BRASILEIRA DE
Entomologia
A Journal on Insect Diversity and Evolutionwww.rbentomologia.com

Systematics, Morphology and Biogeography

Larva, pupa and DNA barcodes of the Neotropical geometrid moth
Glena mielkei (Lepidoptera: Geometridae: Ennominae: Boarmiini)Felipe Méndez-Abarca^a, Sebastián Espinoza-Donoso^b, Scott Escobar-Suárez^b,
Wilson Huanca-Mamani^c, Héctor A. Vargas^b, Enrique A. Mundaca^{d,*}^a Fundación Reino Animal, Cienfuegos S/N, Arica, Chile^b Universidad de Tarapacá, Facultad de Ciencias Agronómicas, Departamento de Recursos Ambientales, Casilla 6-D, Arica, Chile^c Universidad de Tarapacá, Facultad de Ciencias Agronómicas, Departamento de Producción Agrícola, Casilla 6-D, Arica, Chile^d Universidad Católica del Maule, Facultad de Ciencias Agrarias y Forestales, Escuela de Agronomía, Casilla 7-D, Curicó, Chile

ARTICLE INFO

Article history:

Received 19 March 2018

Accepted 24 July 2018

Available online xxx

Associate Editor: Livia Pinheiro

Keywords:

Atacama Desert

*Glena interpunctata**Physocleora*

Immature stages

Trixis cacalioides

ABSTRACT

Glena mielkei Vargas, 2010 (Lepidoptera: Geometridae: Ennominae: Boarmiini) is a Neotropical geometrid moth native to the Atacama Desert of northern Chile whose larvae are folivorous on the shrub *Trixis cacalioides* (Asteraceae). The last instar and pupa are described and illustrated, and DNA barcode sequences are provided for the first time for *G. mielkei*. Descriptions are made based on larvae collected in the type locality. Comparisons with the available descriptions of congeneric species suggest that the chaetotaxy of the SV group of the abdominal segment and the morphology of the cremaster could be useful tools to species identification based on last instar and pupa, respectively. A search in BOLD (Barcode of Life Data System) showed that the only DNA barcode haplotype found in the two specimens sequenced was closest to *Physocleora* Warren, 1897 than *Glena* Hulst, 1896. These results coincide with the morphological peculiarities of the genitalia highlighted in the original description of *G. mielkei*, suggesting that a definitive assessment of the generic status of this geometrid moth deserves further integrative studies.

© 2018 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Species of the family Geometridae tend to exhibit a close relationship with the vegetation (Scoble, 1995; Brehm and Fiedler, 2005; Brehm et al., 2005; Bolte, 1990). Records of the Geometridae fauna in Chile show more than 450 species for the central and southern areas of the country, with most of the described species represented by the subfamilies Ennominae and Larentiinae (Parra, 1995). Records for the north area of Chile, however, remain scarce, with only few species having been described (Vargas, 2010).

Glena Hulst, 1896 is a New World genus of Boarmiini (Lepidoptera: Geometridae: Ennominae) that comprises more than 40 species described so far, of which over 30 occur in the Neotropics (Pitkin, 2002). Natural history and external morphology of immature stages of *Glena* still remain poorly studied (Rindge, 1965, 1967). Plants of the Aceraceae, Ericaceae, Pinaceae, Rosaceae, Salicaceae

and Tamaricaceae families are recorded as hosts for Nearctic representatives of *Glena*, while the host records for the Neotropical fauna include plants of the Asteraceae, Clusiaceae, Cupressaceae, Erythroxylaceae, Fabaceae, Myrtaceae and Pinaceae families (Osorio, 2005; Marconato et al., 2008; Robinson et al., 2010; Méndez-Abarca et al., 2014). However, these records are mostly based on a few polyphagous species, whereas many others have never been reared from larvae.

Glena mielkei Vargas, 2010 is the only species of the genus currently described for Chile. To date, the literature on *G. mielkei* includes a description based on the adult stages plus information about its distributional range, which has so far been restricted to the province of Arica (Northern Chile) (Fig. 1A). Although in the laboratory, larvae of *G. mielkei* are able to feed on three species of Asteraceae, namely *Trixis cacalioides* (Kunth), *Pluchea chingollo* (Kunth) and *Tessaria absinthioides* (Hook. & Arn.), the only host plant recorded in the field has been *T. cacalioides* (Vargas, 2010; Méndez-Abarca et al., 2014) (Fig. 1B). We describe and illustrate the external morphology of the last instar larva and pupa of *G. mielkei* for the first time based on specimens collected in the type locality with the

* Corresponding author.

E-mail: emundaca@ucm.cl (E.A. Mundaca).<https://doi.org/10.1016/j.rbe.2018.07.004>0085-5626/© 2018 Sociedade Brasileira de Entomologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1. Details of the natural history of *G. mielkei* in the Atacama Desert of northern Chile. (A) Type locality in the Azapa Valley, Arica Province, northern Chile. (B) Host plant *T. cacalioides* (Asteraceae) at the type locality. (C) Last instar larva of *G. mielkei* feeding of *T. cacalioides* at the type locality.

aim of contributing to further comparative studies on morphology of immature stages of Boarmiini. In addition, the first DNA barcode (sensu Hebert et al., 2003) sequences of *G. mielkei* are provided.

Materials and methods

Larvae of *G. mielkei* were collected on *T. cacalioides* in the Azapa Valley (type locality) in June 2017. Larvae were placed in plastic vials with leaves of the host plant and towel paper at the bottom and brought to the laboratory. The vials were cleaned and leaves of *T. cacalioides* were provided periodically until larvae finished feeding. Eight larvae (last instar) and six pupae were kept in ethanol 70% to carry out the morphological analysis. The integument of the larvae was cleaned in hot KOH 10% for a few minutes and structures were slide mounted either on glycerine or Euparal to observe morphological details using a Leica M125 stereomicroscope and an Olympus® BX51 optical microscope. Two pupae were kept in ethanol 95% at -20°C until DNA extraction. Four pupae were kept in the plastic vials to obtain adults to confirm the taxonomic identification based on the morphology of the genitalia.

Genomic DNA was extracted from two pupae of *G. mielkei* following the procedures described in Huanca-Mamani et al. (2015). Amplification and sequencing of the DNA barcode region (sensu Hebert et al., 2003) were undertaken at Macrogen, Inc. (South Korea) with the primers LCO-1490 and HCO-2198 (Folmer et al., 1994) following the amplification programme described by Escobar-Suárez et al. (2017). The sequence alignment was performed by the Clustal W method in the software MEGA6 (Tamura et al., 2013), and a search for close sequences was performed in BOLD (Ratnasingham and Hebert, 2007).

Results

Last instar larva (Figs. 1C, 2A–E, 3A–F).

Head well-developed, hypognathous, brown-greyish with irregular darker stains; thorax and abdomen brown-greenish, with two tuberiform dorsolateral projections on A2; prolegs present on A6 and A10 (Fig. 1C).

Head. Two groups of six subcircular stemmata located dorsally from the antennal socket (Fig. 2A and B). Antennae trisegmented: first segment ring-like; second segment cylindrical, width similar to first segment, length about twice the width, pore and sensillum on lateral surface, four sensilla on distal surface; third segment cylindrical, length about one-third that of second segment, width slightly smaller than length. Mouthparts adapted to chew. Labrum bilobulated with straight dorsal margin. Sides slightly convex. Ventral margin with a central cleft. External surface sclerotized, slightly convex with 12 setae and 4 pores. Internal surface slightly concave, membranous with two pores located in the central area and six sclerotized flattened projections by the ventral margin (Fig. 3B and C). Mandibles (Fig. 3D) strongly sclerotized with a pair of bristles nearby the basal area on the external surface (M1, M2), apical margin serrated with seven teeth-like projections. Maxilla (Fig. 3E) with galea and palp well-differentiated; maxillary palp trisegmented; third segment with digitiform sensillum and two pores laterally, distal surface with eight sensilla; galea with six sensilla on distal surface. Labium (Fig. 3F) with cylindrical spinneret located on the apex; labial palpi approximately half the spinneret length.

Thorax (Fig. 2C). Three segments clearly differentiated. Prothorax with a dorsal plate slightly sclerotized and divided along the medial line. Two ellipsoidal spiracles present laterally on prothorax, with conspicuous filtering structures.

Legs (Fig. 2E). Coxa wide, mainly membranous, with eight bristles, three of them very reduced and the remaining five elongated. Trochanter reduced to a slim strip compressed between the coxa and the femur, with three pores and only one highly reduced bristle. Femur cylindrical, with most of its surface highly sclerotized, membranous towards its medial area, with two bristles.

Download English Version:

<https://daneshyari.com/en/article/8953078>

Download Persian Version:

<https://daneshyari.com/article/8953078>

[Daneshyari.com](https://daneshyari.com)