

Accepted Manuscript

Title: A Pilot study using Flash Continuous Glucose Monitoring in Patients with Type-2 Diabetes on Multiple Anti-diabetic Agents During Ramadan

Authors: Mohammed Bashir, Tarik Elhadd, Hamda Ali, Khaled Baagar, Ibrahim A. Abdel Hakam, Dabia H.S. Al-Mohanadi, Emad Naem, Amina Al Abdulla, Kawsar Mohammed, Abdul-Badie Abou-Samra



PII: S1871-4021(18)30210-8
DOI: <https://doi.org/10.1016/j.dsx.2018.06.005>
Reference: DSX 1015

To appear in: *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*

Please cite this article as: Bashir M, Elhadd T, Ali H, Baagar K, Abdel Hakam IA, Al-Mohanadi DHS, Naem E, Al Abdulla A, Mohammed K, Abou-Samra A-Badie, A Pilot study using Flash Continuous Glucose Monitoring in Patients with Type-2 Diabetes on Multiple Anti-diabetic Agents During Ramadan, *Diabetes and Metabolic Syndrome: Clinical Research and Reviews* (2018), <https://doi.org/10.1016/j.dsx.2018.06.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A Pilot study using Flash Continuous Glucose Monitoring in Patients with Type-2 Diabetes on Multiple Anti-diabetic Agents During Ramadan.

Mohammed Bashir, Tarik Elhadd, Hamda Ali, Khaled Baagar, Ibrahim A Abdel Hakam, Dabia H S Al-Mohanadi, Emad Naem, Amina Al Abdulla, Kawsar Mohammed & Abdul-Badie Abou-Samra.

Qatar Metabolic Institute, Department of Medicine, Hamad Medical Corporation;

Doha, Qatar

Address for Correspondence: Dr Mohammed Bashir

Qatar Metabolic Institute, Department of Medicine, Hamad Medical Corporation;

Doha,

Qatar

P O Box 3050

Email:- mohdbashir80@hotmail.com

Abstract

Background: - This study aims to examine the incidence of hypoglycaemia, based on activity, during Ramadan in patients with type 2 diabetes mellitus who were on were on three or more anti-diabetic medications.

Methods: - Type 2 diabetes patients who fasted during Ramadan and were on three or more anti-diabetic medications were studied for two weeks using flash glucose monitoring.

Download English Version:

<https://daneshyari.com/en/article/8957031>

Download Persian Version:

<https://daneshyari.com/article/8957031>

[Daneshyari.com](https://daneshyari.com)