

EDITORIAL COMMENT

# Internal Bleeding

## Is Intraplaque Hemorrhage a Decoration or a Driver?\*



Jean-Baptiste Michel, MD, PhD,<sup>a</sup> Peter Libby, MD,<sup>b</sup> Grégory Franck, PhD<sup>a</sup>

Contemporary advances in high-throughput screening of biological samples have added unbiased screens to the classical candidate approach for implicating mediators in various disease states. In this issue of *JACC: Basic to Translational Science*, Matic et al. (1) report transcriptomic and proteomic profiling of atherosclerotic plaques banked from carotid endarterectomies. In parallel, they looked at blood sampled near the lesion in vivo in comparison to peripheral blood. The investigators found a single molecule of many tested to be overexpressed in this collection of atherosclerotic lesions that received surgical remediation compared with nonatherosclerotic arterial specimens. They identified biliverdin reductase B (BLVRB) as significantly increased at a messenger RNA, local protein, and plasma protein levels, in company with parallel rises in heme oxygenase 1 (HMOX1). BLVRB participates in catabolism of heme.

SEE PAGE 464

These results that emerged from high-throughput screening technologies applied to human atherosclerosis draw our attention to pathways long implicated


in atherogenesis, but remain worthy of renewed attention in light of new data including these from the Stockholm/Karolinska group. The strong regulation of heme-metabolizing enzymes in active human atheromata indicates a substantial local and/or systemic tissue response to heme, at least in carotid artery lesions sufficiently active to warrant surgical treatment.

From whence does heme that could evoke a clearance response within the plaque come? The microvessels that penetrate into plaques during their evolution (inward neo-angiogenesis) exhibit impaired barrier function and display characteristics of fragility that could lead to extravasation of erythrocytes and intraplaque hemorrhage (Figure 1). In the extreme, leaky microvessels in the plaque could give rise to rapid plaque expansion and promote hematoma formation that could embarrass blood flow. Such macroscopic episodes of intraplaque hemorrhage, although uncommon, likely represent the extreme case of much more common microscopic episodes of intraplaque hemorrhage. Matic et al. (1) found that exposure to ferric iron was associated with iron uptake by THP-1 human monocyte cell line-derived macrophage-like cells and stimulated HMOX1 and BLVRB expression, suggesting that local hemolysis and iron derived from heme drive the elevated concentrations of these enzymes in plaques. Intraplaque hemorrhage promotes erythrocyte sequestration and localized hemolysis, hemostasis, and coagulation, including platelet activation, fibrin formation, and trapping of leukocytes, mainly neutrophils, in human plaques (2). Foci of intraplaque hemorrhage colocalize with proteases and markers of oxidative stress, which are possible promoters of plaque rupture (3,4). In contrast, products of local blood coagulation can lead to the release of platelet products, such as transforming growth factor-beta and platelet-derived growth factor, which can foster a local healing response that could augment smooth muscle production of extracellular matrix,

\*Editorials published in *JACC: Basic to Translational Science* reflect the views of the authors and do not necessarily represent the views of *JACC: Basic to Translational Science* or the American College of Cardiology.

From the <sup>a</sup>Unité mixte de Recherche 1148 Inserm-Paris7 University, Xavier Bichat Hospital, Paris, France; and the <sup>b</sup>Division of Cardiovascular Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts. Dr. Libby was supported by grants from the National Heart, Lung, and Blood Institute (HL080472) and the RRM Charitable Fund. Dr. Franck was supported by the Lefoulon-Delalande Foundation and the UE PRESTIGE Fund. Dr. Michel has reported that he has no relationships relevant to the contents of this paper to disclose.

All authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the *JACC: Basic to Translational Science* [author instructions page](#).



smooth muscle proliferation, and thus, growth of the atherosclerotic lesion.

In addition to erythrocyte extravasation from fragile microvessels, entry of erythrocytes from the macrovascular compartment into the intima could arise from breaches in the integrity of the endothelial layer that lines the macrovascular lumen. Altered hydrodynamics at regions of flow disturbance could contribute to endothelial cell desquamation or impaired barrier function. Recent studies suggest accumulation of erythrocytes near the intimal interface with blood even in early atherogenesis (5). Not only do red blood cells (RBCs) accumulate locally within the intima, but hemoglobin, glycophorin A, and iron colocalize with regions of RBC collections that compose small intraplaque hematomas. The sites of local RBC accumulation in the intima also contain HMOX1, ferroportin, and members of the natural resistance-associated macrophage protein family of metal transporters. Furthermore, sites of these intimal hematomas contain markers of oxidation of lipids and proteins. Smooth muscle cells appear capable of erythrophagocytosis in addition to classical scavenging of modified or senescent RBC via scavenger receptors associated with mononuclear phagocytes including CD163 and CD36 (5).

Whether they are derived from microvascular leaks or from breaches in the macrovascular endothelium, redox-active ferrous iron ( $\text{Fe}^{++}$ ) derived from erythrocyte heme prosthetic groups can contribute to local oxidative damage through the Fenton reaction. The local plaque environment favors hemolysis of RBC, releasing heme and iron in the intimal lesions. Recent systems biology analyses point to hydroxyl radical derived through Fenton chemistry as an agonist in atherogenesis (6). In addition, genome-wide association studies have implicated iron metabolism and hepcidin as risk factors for biomarkers of atherosclerosis such as increased intima-media thickness (7). Experimental studies have implicated hepcidin in activating plaque macrophages following erythrophagocytosis. Some experimental data suggest that clearance of hemoglobin by CD163-positive cells and concomitant HMOX1 expression can provide an anti-inflammatory response that can defend against some of the potentially adverse effects of local heme metabolism and accumulation of iron (8). Beyond the role of RBC-derived heme and iron in potentiating atherosclerosis, the membranes of erythrocytes that enter the intima can provide a local source of cholesterol (9).

Download English Version:

<https://daneshyari.com/en/article/8957138>

Download Persian Version:

<https://daneshyari.com/article/8957138>

[Daneshyari.com](https://daneshyari.com)