Title: Field-grown soybean induces jasmonates and defensive compounds in response to thrips feeding and solar UV-B radiation

Authors: Francisco M. Dillon, Hugo D. Chludil, Michael Reichelt, Axel Mithöfer, Jorge A. Zavala

PII: S0098-8472(18)30893-1
DOI: https://doi.org/10.1016/j.envexpbot.2018.08.022
Reference: EEB 3548

To appear in: *Environmental and Experimental Botany*

Received date: 13-6-2018
Revised date: 4-8-2018
Accepted date: 17-8-2018

Please cite this article as: Dillon FM, Chludil HD, Reichelt M, Mithöfer A, Zavala JA, Field-grown soybean induces jasmonates and defensive compounds in response to thrips feeding and solar UV-B radiation, *Environmental and Experimental Botany* (2018), https://doi.org/10.1016/j.envexpbot.2018.08.022

Field-grown soybean induces jasmonates and defensive compounds in response to thrips feeding and solar UV-B radiation.

FRANCISCO M. DILLON[1,3], HUGO D. CHLUDIL[2], MICHAEL REICHELT[4], AXEL MITHÖFER[5], JORGE A. ZAVALA[1,3]

[1] Universidad de Buenos Aires. Facultad de Agronomía. Cátedra de Bioquímica, Avenida San Martín 4453, C1417DSE Buenos Aires, Argentina

[2] Universidad de Buenos Aires. Facultad de Agronomía. Cátedra de Química de Biomoléculas. Avenida San Martín 4453, C1417DSE Buenos Aires, Argentina

[3] INBA/CONICET Avenida San Martín 4453, C1417DSE Buenos Aires, Argentina

[4] Max Planck Institute for Chemical Ecology, Department of Biochemistry, Jena, Germany

[5] Max Planck Institute for Chemical Ecology, Department of Bioorganic Chemistry, Jena, Germany

Correspondence: J. Zavala. E-mail: zavala@agro.uba.ar

Highlights

- Thrips damage in field-grown soybean induced jasmonates and oxo-phytodienoic acid.
- Thrips damage induced an unknown phenolic compound and TPI activity.