

Accepted Manuscript

An NMR strategy to detect conformational differences in a protein complexed with highly analogous inhibitors in solution


John D. Persons, Shahid N. Khan, Rieko Ishima

PII: S1046-2023(18)30014-8
DOI: <https://doi.org/10.1016/j.ymeth.2018.04.005>
Reference: YMETH 4442

To appear in: *Methods*

Received Date: 24 February 2018
Revised Date: 6 April 2018
Accepted Date: 8 April 2018

Please cite this article as: J.D. Persons, S.N. Khan, R. Ishima, An NMR strategy to detect conformational differences in a protein complexed with highly analogous inhibitors in solution, *Methods* (2018), doi: <https://doi.org/10.1016/j.ymeth.2018.04.005>



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

An NMR strategy to detect conformational differences in a protein complexed with highly analogous inhibitors in solution

John D. Persons[‡], Shahid N. Khan[‡], and Rieko Ishima^{*}

Department of Structural Biology, University of Pittsburgh School of Medicine, Biomedical Science Tower 3, 3501 Fifth Avenue, Pittsburgh, PA 15260, USA

[‡]These authors contributed equally.

^{*} Corresponding authors:

Rieko Ishima, Room 1037, Biomedical Science Tower 3, 3501 Fifth Avenue, Pittsburgh, Pennsylvania 15260; Tel: 412-648-9056; Fax: 412-648-9008; Email: ishima@pitt.edu

Key words: NMR, water, relaxation, protein, HSQC, NOE, deuteration

Download English Version:

<https://daneshyari.com/en/article/8961927>

Download Persian Version:

<https://daneshyari.com/article/8961927>

[Daneshyari.com](https://daneshyari.com)