Accepted Manuscript



Improvement of the resistance against early *Mycobacterium tuberculosis*-infection in the absence of PI3Kγ enzyme is associated with increase of CD4+IL-17+ cells and neutrophils

M.P. Cavalcanti-Neto, R.Q. Prado, A.R. Piñeros, C.A. Sérgio, T.B. Bertolini, A.F. Gembre, S.G. Ramos, V.L. Bonato

PII: S1472-9792(18)30054-4

DOI: 10.1016/j.tube.2018.08.009

Reference: YTUBE 1743

To appear in: *Tuberculosis*

Received Date: 9 February 2018

Revised Date: 19 August 2018

Accepted Date: 21 August 2018

Please cite this article as: Cavalcanti-Neto MP, Prado RQ, Piñeros AR, Sérgio CA, Bertolini TB, Gembre AF, Ramos SG, Bonato VL, Improvement of the resistance against early *Mycobacterium tuberculosis*-infection in the absence of PI3Kγ enzyme is associated with increase of CD4+IL-17+ cells and neutrophils, *Tuberculosis* (2018), doi: 10.1016/j.tube.2018.08.009.

Improvement of the resistance against early Mycobacterium tuberculosis-infection in the absence of PI3Kγ enzyme is associated with increase of CD4+IL-17+ cells and neutrophils

Cavalcanti-Neto, MP [a, d], Prado, RQ [a], Piñeros AR [a], Sérgio, CA[a], Bertolini TB [a], Gembre, AF [a], Ramos SG [c], Bonato VL [a, b,*]

[a] Basic and Applied Immunology Programe, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil

[b] Department of Biochemistry and Immunology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil.

[c] Department of Pathology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil.

[d] Laboratory of Cell Signaling and Metabolic Modulation, Institute of Health and Biotechnology, Federal University of Amazonas, Coari, Brazil.

* Corresponding author. Faculdade de Medicina de Ribeirão Preto, Departamento de Bioquímica e Imunologia. Avenida Bandeirantes, 3900 Ribeirão Preto, São Paulo, Brasil. 14049-900

E-mail address: vlbonato@fmrp.usp.br