

Author's Accepted Manuscript

Early carotid artery stenting after onset neurologic symptoms

Gianmarco de Donato, Francesco Setacci, Edoardo Pasqui, Domenico Benevento, Giancarlo Palasciano, Antonio Sterpetti, Luca di Marzo, Carlo Setacci



www.elsevier.com/locate/bios

PII: S0895-7967(18)30001-2
DOI: <https://doi.org/10.1053/j.semvascsurg.2018.02.001>
Reference: YSVAS50545

To appear in: *Seminars in Vascular Surgery*

Cite this article as: Gianmarco de Donato, Francesco Setacci, Edoardo Pasqui, Domenico Benevento, Giancarlo Palasciano, Antonio Sterpetti, Luca di Marzo and Carlo Setacci, Early carotid artery stenting after onset neurologic symptoms, *Seminars in Vascular Surgery*, doi:10.1053/j.semvascsurg.2018.02.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Seminars in Vascular Surgery Issue

Update on Carotid Stent Angioplasty - Indications and Controversies

Guest Editors: Carlo Setacci MD and Antonio Sterpetti MD

Early carotid artery stenting after onset neurologic symptoms

Gianmarco de Donato¹, Francesco Setacci², Edoardo Pasqui¹, Domenico Benevento¹, Giancarlo Palasciano¹, Antonio Sterpetti³, Luca di Marzo³, Carlo Setacci¹.

Institution

1. Vascular Surgery - University of Siena, Italy
2. Vascular Surgery – Casa di Cura Giovanni XXIII, Monastier di Treviso, Italy
3. Vascular Surgery - University Roma La Sapienza, Italy

Corresponding author:

Gianmarco de Donato
Department of Surgery
Unit of Vascular and Endovascular Surgery
University of Siena
Viale Bracci, 53100 Siena – Italy
Tel. +39 0577 585123
Fax +39 0577 585123
E-mail: setacci@unisi.it

Abstract

Multicenter clinical trials level 1 evidence favors the application of carotid endarterectomy (CEA) in symptomatic patients especially the in the elderly cohort. Carotid artery stenting (CAS) has been proposed as a possible early alternative in selected patients after onset of ipsilateral neurologic symptoms. It is well known that the treatment of acute stroke is time-dependent, in patients with acute ischemic stroke

Download English Version:

<https://daneshyari.com/en/article/8963265>

Download Persian Version:

<https://daneshyari.com/article/8963265>

[Daneshyari.com](https://daneshyari.com)