


Accepted Manuscript

Clinical spectrum, treatment and outcome of children with suspected diagnosis of Chronic Inflammatory Demyelinating Polyradiculoneuropathy

A. Silwal , M. Pitt , R. Phadke , K. Mankad , J.E. Davison ,
A. Rossor , C. DeVile , M.M. Reilly , A. Manzur , F. Muntoni ,
P. Munot

PII: S0960-8966(17)31558-4
DOI: [10.1016/j.nmd.2018.06.001](https://doi.org/10.1016/j.nmd.2018.06.001)
Reference: NMD 3560



To appear in: *Neuromuscular Disorders*

Received date: 29 December 2017
Revised date: 2 June 2018
Accepted date: 4 June 2018

Please cite this article as: A. Silwal , M. Pitt , R. Phadke , K. Mankad , J.E. Davison , A. Rossor , C. DeVile , M.M. Reilly , A. Manzur , F. Muntoni , P. Munot , Clinical spectrum, treatment and outcome of children with suspected diagnosis of Chronic Inflammatory Demyelinating Polyradiculoneuropathy , *Neuromuscular Disorders* (2018), doi: [10.1016/j.nmd.2018.06.001](https://doi.org/10.1016/j.nmd.2018.06.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights:

- The diagnosis of CIDP can be challenging.
- In our cohort 52% were diagnosed as CIDP on re-evaluation.
- Cranial nerve abnormality is rare and may be only presenting symptom.
- Children require long-term follow up as the course may be protracted.
- With early treatment majority have good recovery and maintain ambulation.

Download English Version:

<https://daneshyari.com/en/article/8963299>

Download Persian Version:

<https://daneshyari.com/article/8963299>

[Daneshyari.com](https://daneshyari.com)